

A new leucosiid crab (Decapoda) from the middle Pleistocene Atsumi Group, Japan

Hiroaki Karasawa

Mizunami Fossil Museum, Yamanouchi, Akeyo, Mizunami, Gifu 509-6132 Japan <GHA06103@nifty.com>

Abstract

Arcania kobayashinobuakii new species, a leucosiid crab is described from the middle Pleistocene Atsumi Group, central Honshu, Japan. The new species is similar to the extant *Arcania erinacea* (Fabricius, 1787), but differs in arrangement of spines on the carapace.

Key words: Crustacea, Decapoda, Leucosiidae, Pleistocene, Japan

Introduction

The genus *Arcania* Leach, 1817, comprises 20 species from the extant Indo-West Pacific regions (Ng *et al.*, 2008). Five species are recognized as fossils: *Arcania cornuta* (MacGilchrist, 1905) from the Pliocene of Java (Van Straelen, 1938), *A. sp. cfr. A. globata* Stimpson, 1858, from the Pleistocene of Japan (Kato and Karasawa, 1998), *A. elongata* Yokoya, 1933, from the Pleistocene of Japan (Kobayashi *et al.*, 2007), *A. shuishuni* (Hu and Tao, 1985) from the Pliocene of Taiwan; *A. undecimspinosa* De Haan, 1841, from the Pliocene of Taiwan (Hu and Tao, 2000); *A. sp. cf. A. undecimspinosa* De Haan, 1841, from the Pleistocene of Japan (Kato and Karasawa, 1998). Most recently, Vega *et al.* (2010) reported an unnamed species of *Arcania* from the Miocene deposit of Iran.

The purpose of the present paper is to describe a new species of *Arcania* from the middle Pleistocene Atsumi Group, Japan. The specimen was collected from sandy-silt of the Toyohashi Formation of the Atsumi Group exposed at Takamatsu, Tahara City, Aichi Prefecture, central Japan (Karasawa and Tanaka, 1994, Fig. 1). The formation contains rich marine fossils such as mollusks, barnacles, and decapods. Nakashima *et al.* (2008) showed that the Toyohashi Formation was correlated with Marine Isotope Stage 9 (middle Pleistocene).

The specimen described here is deposited in the Mizunami Fossil Museum (MFM).

Systematics

Family Leucosiidae Samouelle, 1819
Subfamily Ebaliinae Stimpson, 1871
Genus *Arcania* Leach, 1817



Fig. 1. *Arcania kobayashinobuakii* new species, MFM 142023 (holotype), carapace, dorsal view, scale bar=5 mm.

Arcania kobayashinobuakii new species

(Fig. 1)

Diagnosis: Carapace strongly convex longitudinally and transversely; dorsal surface spinose with irregular, well-developed spines. Gastric regions with 5 pairs of simple spine; cardiac region with 2 simple spines transversely arranged; intestinal region with median and posterior spinulose spines, a pair of simple spine present at base of posterior spine; branchial regions with 6 simple spines. Anterolateral margin with 4 simple spines; posterolateral margin with 3 spines; posterior margin with 2 large, spinulose spines.

Derivation of name: In honor to my collaborator, Nobuaki

Kobayashi (Gamagouri, Aichi; Assistance of the Mizunami Fossil Museum), who donated his fossil crab collection.

Description: Carapace small, strongly convex longitudinally and transversely. Front, frontal, and anterior gastric regions and marginal part of anterior carapace not preserved. Dorsal surface spinose with well-developed spines of uneven size; regions poorly defined, gastric and intestinal regions separated from branchial regions by shallow groove. Hepatic region swollen with inflated margin fringed with 2 simple spines. Gastric regions ornamented with 5 pairs of simple spine. Cardiac region with 2 simple spines transversely arranged. Intestinal region with median and posterior spinulose spines; posterior spine longest, directed posteriorly; a pair of posterolaterally directed simple spine present at the base of posterior spine. Branchial regions ornamented with 6 simple spines. Anterolateral margin with 2 simple spines behind hepatic region, epibranchial spine not preserved. Posterolateral margin bearing 3 spines; first and third spines simple, second spine spinulose, largest of all spines. Posterior margin bearing 2 large, spinulose spines.

Remarks: The present new species resembles *Arcania erinacea* (Fabricius, 1787) from the Indo-West Pacific (Galil, 2001), but differs in arrangement of spines. The carapace spines of the new species are more sparsely arranged than those of *A. erinacea*.

Material examined: MFM142023 (holotype).

Acknowledgement

I thank N. Kobayashi (Gamagouri, Aichi) for donating his fossil crabs for my study.

References

- De Haan, W. 1833–1850. Crustacea. In, Siebold, P. F. von, Fauna Japonica sive descriptio animalium, quae in Itinere per Japoniam, Jussu et auspiciis superiorum, qui Summun in India Batava Imperium tenent, suscepto, annis 1823–1830 collegit, notis, observationibus et adumbrationibus illustravit. i–xvii+i–xxx+i–x–xvi+1–243, pls. A–J+L–Q+1–55. Ludguni-Batavorum.
- Fabricius, J. C. 1787. Mantissa Insectorum sistens eorum species nuper detectas adiectis Characteribus genericis, Differentiis specificis, Emendationibus, Observationibus 1: i–xx, 1–348. Hafniae.
- Galil, B. S. 2001. A revision of the genus *Arcania* Leach, 1817 (Crustacea: Decapoda: Leucosioidea). *Zoologische Mededelingen* 75(11): 169–206.
- Hu, C.-H., and H.-J. Tao. 1985. Tertiary crustaceans fossil from Taiwan. *Petroleum Geology of Taiwan* 21: 239–259.
- Hu, C.-H., and H.-J. Tao. 2000. Crustacean fossils from southern Taiwan. *Petroleum Geology of Taiwan* 34: 105–195.
- Karasawa, H., and T. Tanaka. 1994. Decapod crustaceans from the Atsumi Group (Middle Pleistocene) of Aichi Prefecture, central Japan. *Science Report of the Toyohashi Museum of Natural History* 4: 11–19.
- Kato, H., and H. Karasawa. 1998. Pleistocene fossil decapod Crustacea from the Boso Peninsula, Japan. *Natural History Research, Special Issue* 5: 1–31.
- Kobayashi, N., T. Goda, N. Ohira, and H. Karasawa. 2004. New records of crabs and barnacles (Crustacea: Decapoda and Cirripedia) from the middle Pleistocene Atsumi Group of Aichi Prefecture, Japan. *Bulletin of the Mizunami Fossil Museum* 34: 111–116.
- Leach, W. E. 1817. *The Zoological Miscellany, being descriptions of new or interesting animals* 3: i–vi, 1–151, pls. 121–149. London.
- MacGilchrist, A. C. 1905. *Natural History Notes from the R.I.M.S.S. "Investigator", Capt. T.H. Heming, R.N. (retired), commanding. Ser III. No. 6. An Account of the new and some of the rarer Decapod Crustacea obtained during the Surveying Seasons 1901–1904.* *Annals and Magazine of Natural History* 16: 361–367.
- Nakashima, R., K. Mizuno, and A. Furusawa. 2008. Depositional age of the Middle Pleistocene Atsumi Group in Atsumi Peninsula, central Japan, based on tephra correlation. *The Journal of the Geological Society of Japan* 114: 70–79.
- Ng, P. K. L., D. Guinot, and P. J. F. Davie. 2008. *Systema Brachyurorum: Part I. An annotated checklist of extant brachyuran crabs of the world.* *Raffles Bulletin of Zoology, Supplement no. 17*: 1–286.
- Samouelle, G. 1819. *The entomologist's useful compendium; or an introduction to the knowledge of British insects, comprising the best means of obtaining and preserving them, and a description of the apparatus generally used; together with the genera of Linné, and the modern method of arranging the classes Crustacea, Myriapoda, Spiders, Mites and Insects, from their affinities and structure, according to the views of Dr. Leach. Also an explanation of the terms used in entomology; a calendar of the times of appearance and usual situations of near 3,000 species of British insects; with instructions for collecting and fitting up objects for the microscope.* London. 496 pp.
- Stimpson, W. 1858. *Prodromus descriptionis animalium evertibratorum, quae in expeditione ad Oceanum Pacificum Septentrionalem, a Reipublica Federata missa, Cadwaladaro Ringgold et Johanne Rodgers Ducibus, observavit et descripsit W. Stimpson. Pars VI. Crustacea Oxystomata.* *Proceedings of the Academy of Natural Sciences of Philadelphia* 10: 158–163.

- Stimpson, W. 1871. Preliminary report on the Crustacea dredged in the Gulf Stream in the Straits of Florida by L.F. de Pourtales, Assist. U. S. Coast Survey. Part I. Brachyura. Bulletin of the Museum of Comparative Zoölogy at Harvard College 2: 109–160.
- Van Straelen, V. 1938. Crustacés Décapodes Cénozoïques des Indes Orientales Néerlandaises. Leidsche Geologische Mededeelingen X(1): 90–103.
- Vega, F. J., H. Gholamalian, and A. Bahrami. 2010. First record of Miocene crustaceans from Hormozgan Province, Southern Iran. *Paläontologische Zeitschrift* 84: 485–493.
- Yokoya, Y. 1933. On the distribution of decapod crustaceans inhabiting the continental shelf around Japan, chiefly based upon the materials collected by S. S. Sôyô-Marû, during the year 1923–1930. *Journal of the College of Agriculture, Tokyo Imperial University* 12: 1–225.

Manuscript accepted on June 30, 2013