On some Paleogene and Neogene crabs of Kachchh, Western India

Jyotsana Rai¹, Vijay Prakash Mishra², Ashok Sahni³, Abha Singh¹ and Francisco J. Vega⁴

 ¹Birbal Sahni Institute of Palaeobotany, Lucknow- 226007
 ²4/490, Vivek Khand Gomti Nagar, Lucknow- 226010
 ³98, Mahatma Gandhi Marg, Lucknow, 226001
 ⁴Instituto de Geología, UNAM, Ciudad Universitaria, Coyoacán, México, DF, 04510,

Abstract

Three species of crabs are here reported from the Lower Miocene Khari Nadi Formation at Matanomadh and Lakhpat sections, Western India. *Philyra karkata* new species is included. *Palaeocarpilius* cf. *P. rugifer* Stoliczka, 1871 is first reported from the Middle Eocene Fulra Limestone at Lakhpat section.

Key words: Decapoda, Brachyura, Cenozoic, India

Introduction

Paleogene and Neogene crabs have been recently reported from Western India by several authors (Schweitzer *et al.*, 2004; Ralte *et al.*, 2009; Vega *et al.*, 2010, among others). We report here occurrence of *Palaeocarpilius* cf. *P. rugifer* Stoliczka, 1871 from the Middle Eocene Fulra Limestone at Lakhpat section. *Raninoides* sp., *Philyra karkata* new species and *Hexapinus*? sp. occur in Lower Miocene rocks of the Khari Nadi Formation at Matanomadh and Lakhpat sections. Although preservation of most specimens is not ideal, this report is important to reinforce paleobiogeographic affinities of Miocene brachyuran assemblages with those of the same age for the Indo-Pacific region.

Geology of the Area

The best developed sequence in the Kachchh basin lies in the southwestern part and is represented by paralic largely Paleogene/Neogene sequences overlying the Deccan basalts. Palaeocene sediments consist of red lateritic and tuffaceous, trap-wash material which contain plant fossils and represent fluvio-lacustrine conditions (Fig.1). Wynne (1872) informally classified the Paleogene/Neogene sediments of Kachchh into seven "groups". The time-stratigraphic classification for these rocks was proposed earlier (Biswas, 1965; 1971; 1972) and later the lithostratigraphic classification was designated (Biswas and Raju, 1973; Biswas, 1992) containing eight formations with type sections and details of contacts. The formations in descending order are as follows:

- 1. Matanomarh Formation = ?Palaeocene
- 2. Naredi Formation = Lower Eocene
- 3. Harudi Formation = Late Middle Eocene
- 4. Fulra Limestone Formation = Late Middle Eocene
- 5. Maniara Fort Formation = Oligocene
- 6. Khari Nadi Formation = Lower Miocene
- 7. Chhasra Formation = Upper Miocene
- 8. Sandhan Formation = Pliocene

The earliest Paleogene marine transgression is represented by the Naredi Formation. From its basal succession a diversified datable dinoflagellate assemblage assignable to the Lower Ypresian has been recovered (Garg et al. 2011). The Harudi and Fulra Limestone formations contain datable calcareous nannofossils, dinoflagellates planktonic and larger foraminfers and a precise Bartonain age is suggested for these formations (Rai, 1997; 2007). Late Eocene (Priabonian) is a hiatus in entire Kachchh Basin. The Maniara Fort and Vinjhan Shale formations also contain datable calcareous nannofossils. The Chhasra Formation is represented by the lower Claystone Member with grey and khaki coloured, laminated gypseous shales with alternating thin hard yellowish highly fossiliferous limestones. The upper Member is mainty represented by siltstone with alternating micaceous siltstone and laminated silty shales of khaki colour. In southwestern Kutch, Mesozoic and the Paleogene/Neogene sequences are covered by recent and sub-recent alluvium.

The giant fossil crab *Palaeocarpilius* cf. *P. rugifer* was recovered from the Fulra Limestone Formation present near Lakhpat Fort cliff section (68°46'30.43"E : 23°49'52.57"N), whereas all other small fossil crabs were obtained from the Miocene Maniara Fort Formation exposed on Matanomarh - Denma Road (68°57'E : 23°32'37"N) and near Lakhpat (68°47'E : 23° 49'30"N). Additionally, the Miocene crab bearing horizons are found associated with mainly coastal dwelling vertebrates such as fishes, turtles, crocodiles and sea cows (Sahni and Mishra, 1975) shown in Fig. 2.

Systematic Paleontology

Section Raninoida Ahyong, Lai, Sharkey, Colgan and Ng, 2007 Superfamily Raninoidea De Haan, 1839 Family Raninidae De Haan, 1839 Subfamily Raninoidinae Lőrenthey, in Lőrenthey and Beurlen, 1929 Genus *Raninoides* H. Milne Edwards, 1837 *Type species: Raninoides laevis* Latreille, 1825, by monotypy.



Fig. 1. Geological map of North Western Kachchh indicating formational boundaries and sample locations.

Raninoides sp.

(Pl. 1, Figs. 1, 2)

Description of material: Carapace small, rectangular, elongate, longitudinally nearly flat; orbitofrontal margin about three-fourths the maximum carapace width, small fissure on lateral portion; anterolateral margin short, with small, short spine directed forwards; posterolateral margin long, convex, about four-fifths maximum carapace length; posterior margin curved, two-thirds maximum carapace width; cervical groove sinuous, relatively deep; deep, curved gastrocardiac grooves at midlength of carapace; median ridge extends to anterior margin from middle portion of carapace.

Locality and repository: Lower Miocene Khari Nadi Formation from Lakhpat and Matanomadh Geology Department, Lucknow University, Lucknow. Specimens L.U.V.P 11601; L.U.V.P. 11602.

Measurements (in mm): Specimen L.U.V.P 11601, carapace length = 13.4; width = 8.7; specimen L.U.V.P 11602, carapace length = 15.4, width = 9.8.

Remarks: The two specimens are too incomplete to offer an identification at species level, but are similar to *Raninoides morrisi* Collins, Lee and Noad, 2003, from the Miocene of Sabah. There are also

similarities with the specimen of *Raninoides* sp. figured by Karasawa *et al.* (2008) from the Pleistocene of the Philippines.

Section Eubrachyura de Saint Laurent, 1980 Subsection Heterotremata Guinot, 1977 Superfamily Leucosioidea Samouelle, 1819 Family Leucosiidae Samouelle, 1819 Subfamily Ebaliinae Stimpson, 1871 Genus *Philyra* Leach, 1817

Type species: Leucosia globus Fabricius, 1775, by subsequent designation of H. Milne-Edwards (1837).

Philyra karkata new species (Pl. 1, Figs. 3–8)

Diagnosis: Carapace subovate; anterolateral margin inclined; posterolateral margin broadly rounded; posterior margin straight; anterior surface of carapace smooth; dorsal surface covered by granules; fused urogastric and cardiac regions; female abdomen cavity broadly rounded.

Description: Carapace longitudinally subovate; front narrow; anterolateral margin inclined, rimmed with small, evenly-spaced



Fig. 2. Stratigraphic sections of study areas indicating occurrence of fossiliferous horizons.

tubercles; posterolateral margin broadly rounded; posterior margin nearly straight, about half the maximum carapace width; anterior surface of carapace smooth; dorsal surface covered by uniformly-sized, evenlyspaced coarse granules; deep, posterolaterally directed groove separetes fused urogastric and cardiac regions from branchial areas; base of cardiac region rounded; female abdomen cavity broadly rounded; outer edge of sternites granulose.

Etymology: Species is named after Sanskrit word for crab, karkata.

Locality and repository: Lower Miocene Khari Nadi Formation from Matanomadh Geology Department, Lucknow University, Lucknow. Holotype, L.U.V.P. 11603, paratypes, L.U.V.P. 11606; L.U.V.P. 11607.

Measurements (in mm): Specimen Number 11603, carapace length = 11.8; width = 10.5; Specimen Number 11606, carapace length = 12.0; width = 9.75; Specimen Number 11607, carapace length = 12.0; width = 10.0.

Remarks: Species assigned to the genus Philyra were reviewed by

Galil (2009), who suggested that Philyra sensu stricto was represented by only two extant species, and present a list of 27 species as Philvra sensu lato, pending for further systematic revision. The Indian specimens cannot be placed on any of the seven genera erected by Galil (2009) to contain species previously assigned to Philyra, since their morphological features are quite different. Karasawa et al. (2008) cited 16 fossil species of Philyra from the Miocene to Pleistocene of the Indo-Pacific region. On their systematic list of fossil Decapoda, Schweitzer et al. (2010) listed 18 species, also from the Indo-Pacific region. Since no fossil species was considered by Galil (2009) on her review of species of Philyra, we follow here the classification presented by Schweitzer et al. (2010). Our specimens differ from Philyra trusanensis Collins, Lee and Noad, 2003 and P. granulosa Morris and Collins, 1991, both from the Miocene of Sarawak, in being more globose, the anterior margin is slightly wider and the posterior margin is narrow and has two posterolateral projections. Philyra sp. cf. P. actidens Chen, 1987, reported from the Miocene of the Philippines (Karasawa et al., 2008) has weaker granules and straight furrows that separate branchial areas from cardiac and urogastric regions. Philyra nishimotoi Karasawa, 1989 and P. plana Karasawa, 1989 from the Miocene of Japan, are smaller, have a more rhombic shape of carapace, broader front and weak dorsal granules. Philyra hayasakai Karasawa and Inoue, 1992 from the Miocene of Japan, has very deep furrows on the dorsal carapace and granules are weaker. Philvra mivamotoi Karasawa and Kishimoto, 1996 from the Miocene of Japan, is more rounded (much wider), has granules on lateral margins and a dorsal surface nearly smooth. Philyra ferrica Hu and Tao, 1996, and P. shihcheni (Hu and Tao, 1979), both from the Miocene of Taiwan are both more globose and granules of the dorsal carapace are weaker and scarce.

Superfamily Carpilioidea Ortmann, 1893 Family Carpiliidae Ortmann, 1893 Genus Palaeocarpilius A. Milne-Edwards, 1862 Type species: Cancer macrocheilus Desmarest, 1822, by subsequent designation (= P. macrocheilus var. coronata Bittner, 1886).

Palaeocarpilius cf. P. rugifer Stoliczka, 1871 (Pl. 1; Figs. 9, 10)

Description: Carapace of medium size, subovate transversely, one-third wider than long, widest at lower third; transverse ridges on posterior third of carapace reach posterior tubercle of anterolateral margin, anterior margin slightly curved; anterolateral margin curved, three-fourths the maximum carapace length, with seven strong rounded tubercles, the three anterior tubercles more rounded and less prominent, remainder tubercles become longer, most posterior tubercle being stronger; posterolateral margin smooth, one-third the maximum carapace width, inclined 45° respect transversal axis. Female abdomen subovate elongated, telson plus somites 6 to 3 reach half the maximum carapace length, width is one-third the maximum carapace width; telson subtriangular, lateral margins broadly rounded, one fifth the maximum carapace length, one-sixth the maximum carapace width; somite 6 rectangular, slightly wider but shorter than telson; somite 5 rectangular, as wide as somite 6 but half its length; somite 4 similar to somite 5 but slightly shorter; somite 3 very similar in size and shape to somite 4; remainder somites not seen. Chelipeds robust; coxae subrectang 0.5 d from Oligocene and Miocene deposits of India (Stoliczka, 1871; Ralte *et al.*, 2009; Vega *et al.*, 2010). The specimen here reported is rather small when compared with the mean size of *P. rugifer*, and it was found on Eocene sediments. For this reason, its identification is preliminary until more and more complete specimens are found. This may extend stratigraphic range for *P. rugifer* from the Eocene to the Miocene in India.

Superfamily Hexapodoidea Miers, 1886 Family Hexapodidae Miers, 1886 Genus *Hexapinus* Manning and Holthius, 1981 *Type species: Hexapus latipes* De Haan, 1835, by original designation.

Hexapinus? sp.

(Pl. 1; Figs. 11-14)

Description: Carapace rectangular, medium-sized, twice wider than long; surface covered with evenly-spaced fine granules; front straight, half the maximum carapace width; anterolateral margin slightly curved; posterolateral margin rimmed; posterior margin straight, twothirds the maximum carapace width; protogastric region defined by subtle elevation; metagastric region poorly defined; urogastric region narrow and short; cardiac region an inverted triangle with inverted triangular, elongate metabranchial lobes; cervical groove sinuous, clearly impressed; branchiocardiac groove not so evident, extends to anterolateral margins; epi and mesobranchial regions separated by a shallow, tangential groove. Sternum subtrapezoidal, sternite 4 also subtrapezoidal, with notches on anterolateral margins; male telson triangular, slightly narrow and curve at anterior margin; other sternites and somites not preserved.

Locality and repository: Lower Miocene Khari Nadi Formation from Lakhpat and Matanomadh Geology Department, Lucknow University, Lucknow. Specimens L.U.V.P. 11604; L.U.V.P. 11605.

Measurements (in mm): Specimen L.U.V.P. 11604, carapace length = 14.8; width = 27.7; specimen L.U.V.P. 11605, carapace length = 11.7, width = 29.1.

Remarks: According to a review of the Hexapodidae (Huang *et al.*, 2002), only three extant species represent the genus *Hexapinus*. Presence of the male abdomen, third maxillipeds and pereiopods is extremely important in this family in order to offer a complete diagnosis. Since none of our specimens preserve appendages, we prefer to leave their identification to the generic level, questionably assigned to *Hexapinus*, based on similarities in carapace regions. *Hexapinus* sp. has been reported from the Miocene of Japan (Takeda *et al.*, 1986; Mizuno and Takeda, 1993).

References

- Ahyong, S. T., J. C. Y. Lai, D. Sharkey, D. J. Colgan, and P. K. L. Ng (2007), Phylogenetics of the brachyuran crabs (Crustacea: Decapoda): the status of Podotremata based on small subunit nuclear ribosomal RNA. Molecular Phylogenetics and Evolution, 45, 576–586.
- Biswas, S. K. (1965), A new classification on the Tertiary rocks of Kutch, Western India. Bulletin of the Geological, Mining and Metallurgical Society of India, 35, 1–6.
- Biswas, S. K. (1971), Note on the Geology of Kutch. Quarterly Journal of the Geological Minning and Metallurgical Society of India, 43(4), 223 –235.
- Biswas, S. K. (1972), Time stratigraphic classification of the Tertiary rocks

Plate 1

43



1, 2. *Raninoides* sp. 1, Dorsal view of specimen L.U.V.P 11601; 2, Dorsal view of specimen L.U.V.P. 11602. 3–8. *Philyra karkata* new species. 3, Dorsal view of holotype, L.U.V.P. 11603; 4, Ventral view of same specimen; 5, Dorsal view of paratype L.U.V.P. 11606; 6, Ventral view of same specimen; 7, Dorsal view of paratype L.U.V.P. 11607; 8, Ventral view of same specimen; 9, 10. *Palaeocarpilius* cf. *P. rugifer* Stoliczka, 1871. 9, Dorsal view of specimen Number 39991, 10, Ventral view of same specimen. 11–14. *Hexapinus?* sp. 11, Dorsal view of specimen L.U.V.P. 11604; 12, Ventral view of same specimen; 13, Dorsal view of specimen L.U.V.P. 11605; 12, 14, Ventral view of same specimen. Scale bars = 1 cm.

of Kutch – A Revision and Amendments. Quarterly Journal of the Geological Mining and Metallurgical Society of India, 44(3), 221 – 224.

- Biswas, S. K. (1992), Tertiary Stratigraphy of Kutch. The Palaeontological Society of India, 37, 1–29.
- Biswas, S. K. and D. S. N. Raju (1973), The rock-stratigraphic classification on the Tertiary sediments of Kutch. Bulletin Oil and natural Gas Commission, 10 (1and 2), 37–45.
- Bittner, A. (1886), Neue Brachyuren des Eozäns von Verona. Sitzungsberichte der kaiser Akademie der Wissenschaften in Wien, Mathematisch-Naturwissenschaftliche Klasse, 94, I abteilung, 44–55.
- Chen, H. (1987), On two new species of Leucosiidae (Crustacea: Brachyura) from the Chinese waters. Studia Marina Sinica, 28, 186–203.
- Collins , J. S. H., C. Lee and J. Noad (2003), Miocene and Pleistocene crabs (Crustacea, Decapoda) from Sabah and Sarawak. Journal of Systematic Palaeontology, 1(3), 187–226.
- Desmarest, A. G. (1822), Histoire Naturelle des Crustacés Fossiles. Les Crustacés Proprement Dits. F. G. Levrault, Paris, pp. 67–154.
- Fabricius, J. C. (1775), Systema entomologiae, sistens insectorum classes, ordines, genera, species, adjectis synonymis, locis, descriptionibus, observationibus. Libraria Kortii, Flensburgi et Lipsiae, 832 p.
- Galil, B. S. (2009), An examination of the genus *Philyra* Leach, 1817 (Crustacea, Decapoda, Leucosiidae) with descriptions of seven new genera and six new species. Zoosystema, 31(2), 279–320.
- Garg, R., V. Prasad, B. Thakur, I. B. Singh and K. Ateequzzaman (2011), Dinoflagellate cysts from the Naredi Formation, Southwestern Kutch, India implications on age and palaeoenvironment. Journal of the Palaeontological Society of India, 56(2), 201–218.
- Guinot, D. (1977), Propositions pour une nouvelle classification des Crustacés Décapodes Brachyoures. Comptes rendus hebdomadaires des séances de l'Académie des sciences, série D, 285, 1049–1052.
- Haan, W. de (1833–1850), Crustacea. In Siebold, P. F. von, Fauna Japonica sive Descriptio Animalium, Quae in Itinere per Japoniam, Jussu et Auspiciis Superiorum, qui Summum in India Batava Imperium Tenent, Suscepto, Annis 1823–1830 Collegit, Noitis, Observationibus et Adumbrationibus Illustravit. Leiden, Lugduni-Batavorum. Pp. i–xvii, i–xxxi, ix–xvi, 1–243.
- Hu, C.-H. and H.-J. Tao (1979), Studies on Quaternary fossil crabs and associated faunas from Hainan island, southern China. Proceedings of the Geological Society of China, 22, 145–158.
- Hu, C.-H. and H.-J. Tao (1996), Crustacean Fossils of Taiwan. San-Ming Book, Taipei, 228 pp.
- Huang, J.-F., P.-W. Hsueh and P. K. L. Ng (2002), Crabs of the Family Hexapodidae (Decapoda: Brachyura) from Taiwan, with description of a new genus and new species. Journal of Crustacean Biology, 22, 651–660.
- Karasawa, H. (1989), Decapod crustaceans from the Miocene Mizunami Group, central Japan. Part 1. Thalassinoidea, Leucosioidea and Grapsidoidea. Bulletin of the Mizunami Fossil Museum, 16, 1–28.
- Karasawa, H. and K. Inoue (1992), Decapod crustaceans from the Miocene Kukinaga Group, Tanegashima Island, Kyushu, Japan. Tertiary Research, 14(2), 73–96.
- Karasawa, H. and S. Kishimto (1996), Two new species of decapods crustaceans from the Katsuta Group (middle Miocene), Japan. Bulletin of the Mizunami Fossil Museum, 23, 39–50.
- Karasawa, H., H. Kato, T. Kase, Y. Maac-Aguilar, Y. Kurihara, H. Hayashi and K. Hagino (2008), Neogene and Quaternary Ghost Shrimps and Crabs (Crustacea: Decapoda) from the Philippines. Bulletin National Museum of Natural Sciences, Series C, 34, 51–76.
- Latreille, P. A. (1825), Familles naturelles du règne animal, exposés succinctement et dans un ordre analytique, avec l'indication de

leurs genres. Bailliere, Paris. 570 pp.

- Leach, W. E. (1817), Monograph on the Genera and Species of the Malacostracous Family Leucosidea. *In*: Leach, W. E., The Zoological Miscellany; being descriptions of new, or interesting animals. Vol. 3. Covent Garden and London, E. Nodder and Son. p. 17–26.
- Lőrenthey, E. and K. Beurlen (1929), Die fossilen Dekapoden der Länder der Ungarischen Krone. Geologica Hungarica, Series Palaeontologica, 3, 421 pp.
- Manning, R. B. and L.B. Holthuis (1981), West African brachyuran crabs (Crustacea: Decapoda). Smithsonian Contributions to Zoology, 306, 1–379.
- Miers, E. J. (1886), Report on the Brachyura collected by H.M.S. Challenger during the years 1873–1876. In: Murray, J., Zoology. Wyville Thomson, C. & J. Murray, Report on the Scientific Results of the Voyage of H.M.S. Challenger During the Years 1873–76 Under the Command of Captain George S. Nares, R.N., F.R.S. and the Late Captain Frank Tourle Thomson, R.N. Vol. 17. Edinburgh, Neill and Company. p. 1–362.
- Milne-Edwards, A. (1862), Monographie des Crustacés Fossiles de la Famille des Canceriens. Annales des Sciences Naturelles, 4e série, 18, 31– 85.
- Milne-Edwards, H. (1834–1840), Histoire naturelle des Crustacés, comprenant l'anatomie, la physiologie et la classification de ces animaux. Librairie Encyclopédique de Roret. Vol. 1–3. Paris, Roret. (1)468, (2)532, (3)638 pp.
- Mizuno, Y. and M. Takeda (1993), Crustacea, p. 77–90, pls. 1–5. In Tokai Fossil Society (ed.), Fossils from the Miocene Morozaki Group, Tokai Fossil Society, Nagoya.
- Morris, S. F. and J. S. H. Collins (1991), Neogene crabs from Brunei, Sabah and Sarawak. Bulletin of the British Museum of Natural History (Geology), 47(1), 1–33.
- Ortmann, A. E. (1893), Die Decapoden-Krebse des Strassburger Museums, mit besonderer Berücksichtigung der von Herrn Dr. Döderlein bei Japan und bei den Liu-Kiu-Inseln gesammelten und zur Zeit im Strassburger Museum aufbewahrten Formen. VII. Theil. Abtheilung: Brachyura (Brachyura genuina Boas) II. Unterabtheilung: Cancroidea, 2. Section: Cancrinea, 1. Gruppe: Cyclometopa. Zoologische Jahrbücher. Abteilung für Systematik, Geographie und Biologie der Thiere, 7(3), 411–495.
- Rai, J. (1997), Scanning electron microscopic studies of the late middle Eocene (Bartonian) calcareous nannofossils from the Kutch Basin, western India. Journal of the Palaeontological Society of India, 42, 147–167.
- Rai, J. (2007), Middle Eocene calcareous nannofossil biostratigraphy and Taxonomy of onland Kutch Basin, western India, The Paleobotanist, 56, 29–116.
- Ralte, V. Z., J. Lalchawimawii, J. Malsawma and R. P. Tiwari (2009), Decapod fossils from the Bhuban Formation, Surma Group, Aizawl, Mizoram. Earth Science India, 2(III), 196–210.
- Sahni, A. and V. P. Mishra (1975), Lower Tertiary vertebrates from Western India, Monograph of the Palaeontological Society of India, 3, 1– 48.
- Saint Laurent, M. de (1980), Sur la classification et la phylogénie des Crustacés Décapodes Brachyoures. I. Podotremata Guinot, 1977 et Eubrachyura sect. nov. Comptes rendus hebdomadaires des séances de l'Académie des sciences, série III, 290, 1265–1268.
- Samouelle, G. (1819), The entomologist's useful compendium; or an introduction to the knowledge of British insects, comprising the best means of obtaining and preserving them, and a description of the apparatus generally used; together with the genera of Linné, and

the modern method of arranging the classes Crustacea, Myriapoda, Spiders, Mites and Insects, from their affinities and structure, according to the views of Dr. Leach. Also an explanation of the terms used in entomology; a calendar of the times of appearance and usual situations of near 3,000 species of British insects; with instructions for collecting and fitting up objects for the microscope. London. 496 pp.

- Schweitzer, C. E., R. M. Feldmann and P. D. Gingerich (2004), New Decapoda (Crustacea) from the Middle and Late Eocene of Pakistan and a revision of *Lobonotus* A. Milne Edwards 1864. Contributions from the Museum of Paleontology, The University of Michigan, 31(4), 89–118.
- Schweitzer, C. E., R. M. Feldmann, A. Garassino, H. Karasawa, and G. Schweigert (2010), Systematic list of fossil decapod crustacean species. Crustaceana Monographs. Fransen, C. H. J. M. and J. C. von Vaupel Klein (eds.) Vol. 10. Leiden: Brill. 230 pp.
- Stimpson, W. (1871), Preliminary report on the Crustacea dredged in the Gulf Stream in the Straits of Florida by L.F. de Pourtales, Assist.

U. S. Coast Survey. Part I. Brachyura. Bulletin of the Museum of Comparative Zoölogy at Harvard College, 2, 109–160.

- Stoliczka, F. (1871), Observations of fossil crabs from Tertiary deposits in Sind and Kutch. Memoirs of the Geological Survey of India, Palaeontologica Indica, series 7, XIV, volume 1, part 1, 16 pp.
- Takeda, M., Y. Mizuno and M. Yamaoka (1986), Some fossil crustaceans from the Miocene Morozaki Group in the Chita Peninsula, central Japan. Kaseki no Tomo, 28, 12–22.
- Vega, F. J., J. K. Tiwari and S. Bajpai (2010), Additions to *Palaeocarpilius rugifer* Stoliczka from the Oligocene of Kutch, western India. Bulletin of the Mizunami Fossil Museum, 36, 45–49.
- Wynne, A. B. (1872), Memoir on the Geology of Kutch, to accompany the map compiled by A.B. Wynne and F. Fedden during the season 1867–68,1868–69. Memoir Geological Survey of India, 19(2), 1– 269.

Manuscript accepted on May 11, 2012