

Axiidea and Brachyura (Decapoda) from the Pliocene–Pleistocene Ananai Formation, Shikoku, Japan

Hiroaki Karasawa¹ and Yoshitaka Kinugawa²

¹Mizunami Fossil Museum, Yamanouchi, Akeyo, Mizunami, Gifu 509-6132, Japan

<GHA06103@nifty.com>

²Kaminotani7-3-3, Suma-ku, Kobe City, 654-0162, Japan

Abstract

Four species of decapods are reported from the Ananai Formation of the Pliocene–Pleistocene Tonohama Group in Aki City, Kochi Prefecture, Shikoku, Japan. *Ctenocheles* sp. (Axiidea: Ctenochelidae), *Anatolikos japonicus* (Ortmann) (Brachyura: Cancridae), and two species of *Charybdis* (Brachyura: Portunidae) are included. *Charybdis (Charybdis) kinugawai* Karasawa, a new portunid is described.

Key words: Decapoda, Axiidea, Brachyura, Ananai Formation, Pliocene–Pleistocene, Japan

Introduction

The Pliocene–Pleistocene Tonohama Group is distributed in the western part of the Muroto Peninsula, Kochi Prefecture, Shikoku. This group consists of the Nobori, Ananai, and Ropponmatsu Formations in ascending order (Katto and Masuda, 1993; Iwai et al., 2006). Matsubara (2004) reviewed the geologic age of the Tonohama Group based upon the previously known microfossils. He showed that the Nobori Formation was assigned to the late early Pliocene–early late Pliocene (4.20–3.21 or 3.12 Ma; Berggren et al., 1995) and that the geological age of the Ananai Formation was the late Pliocene (2.78 or 2.73–1.97 Ma) (=late Pliocene–early Pleistocene based upon Cohen and Gibbard, 2011).

The Tonohama Group contains rich marine fossils (i.e., Iwai et al., 2006). However, decapods have been very rare; Mimoto ed. (1986) recorded three brachyurans, *Charybdis* sp., *Leucosia* sp., and *Parthenope* sp., from the Ananai Formation and, subsequently, Mimoto (2001) described one axiidean Callianassoidea gen. et sp. indet., and four brachyurans, Xanthoidea ? gen. et sp. indet., Leucosiidae gen. et sp. indet., and *Macrocheira* sp., from the Nobori Formation. The junior author (YK) collected seven specimens of decapods from fine-grained sandstone of the Ananai Formation exposed at Tonohama (Loc. 3 of Iwai et al., 2006), Aki City, Kochi Prefecture. The purpose of this work is to describe and figure one axiidean and three brachyurans including a new species of the portunid genus *Charybdis* de Haan, 1833.

The described specimens are housed in the Mizunami Fossil Museum (MFM).

Systematics (By Karasawa)

Infraorder Axiidea de Saint Laurent, 1979

Superfamily Callianassoidea Dana, 1852

Family Ctenochelidae Manning and Felder, 1991

Subfamily Ctenocheliniae Manning and Felder, 1991

Genus *Ctenocheles* Kishinouye, 1926

Ctenocheles sp.

(Fig. 1.1, 1.2)

Material examined: Incomplete propodi of 1st pereiopod, MFM142181, 142182.

Remarks: The Japanese fossil species of *Ctenocheles* comprise *Ctenocheles sujakui* Imaizumi, 1958, from the lower Oligocene Kishima Formation (Imaizumi, 1958; Karasawa, 1993, 1997; Karasawa and Fudouji, 2000) and an unnamed species from the middle Miocene Shimo Formation (Karasawa et al., 2011). *Ctenocheles balssi* Kishinouye, 1926, is only known from the Japanese waters (Sakai, 2010). However, I do not make sufficient comparison because of the poorly preserved material.

Infraorder Brachyura Linnaeus, 1758

Section Eubrachyura de Saint Laurent, 1980

Superfamily Cancroidea Latreille, 1802

Family Cancridae Latreille, 1802

Subfamily Cancrinae Latreille, 1802




Fig. 1, 2. *Ctenocheles* sp., 1, MFM142181, right propodus of 1st cheliped, mesial view; 2, MFM142182, left propodus of 1st cheliped, lateral view. **3.** *Charybdis* sp., MFM142184, carapace, dorsal view. **4.** *Anatolikos japonicus* (Ortmann, 1893), MFM142183, carapace, dorsal view. **5, 6.** *Charybdis (Charybdis) kinugawai* Karasawa, new species, 5, MFM142022 (paratype), carapace, dorsal view; 6. MFM142021 (holotype), carapace, dorsal view. Scale bar = 1 cm.

Genus *Anatolikos* Schweitzer and Feldmann, 2000

***Anatolikos japonicus* (Ortmann, 1893)**

(Fig. 1.4)

Cancer japonicus Ortmann, 1893, p. 427, pl. 17, fig. 5.

Anatolikos japonicus (Ortmann, 1893); Schweitzer and Feldmann, 2000, p. 229.

Material examined: One nearly complete carapace, MFM142183.

Remarks: The extant species ranges from Japan to Taiwan (Sakai, 1976; Ng et al., 2001). The fossil records of the species have been known from the Pleistocene Dainichi Formation of the Kakegawa Group (Karasawa, 1993, 1997) and Pleistocene Onma and Zukawa Formations (Karasawa, 1993; 1997). Hu and Tao (2000) reported the present species from the Pleistocene Ryukyu Group in Taiwan (Hu and Tao, 2000, p. 130, pl. 5, figs. 3, 6); however, the illustrated specimen should be identical with a species of the xanthid *Eitisus* H. Milne Edwards, 1834, in that a smooth carapace has 12 anterolateral spines, the well-defined sterno-abdominal cavity is located on a wide posterior thoracic sternite 4, and the propodus of the cheliped is smooth laterally and ventrally. Therefore, the fossil record of *A. japonicus* is restricted in Japan.

Superfamily Portunoidea Rafinesque, 1815, sensu Karasawa et al., 2008

Family Portunidae Rafinesque, 1815, sensu Karasawa et al., 2008

Subfamily Thalamitinae Paul'son, 1875, sensu Karasawa et al., 2008

Genus *Charybdis* de Haan, 1833

***Charybdis* sp.**

(Figs. 1.3)

Material examined: MFM142184, 142185.

Remarks: Within the present specimen the details of the frontal and anterolateral margins are not known. I do not make sufficient comparison because of the poor-preserved carapaces.

Subgenus *Charybdis* de Haan, 1833

***Charybdis* (*Charybdis*) *kinugawai* Karasawa, new species**

(Figs. 1.5, 1.6; Fig. 2)

Material examined: MFM142021 (holotype), 142022 (paratype).

Diagnosis: Carapace wider than long, length about 75 percent carapace width. Front with 8 teeth; median frontal teeth broadly

triangular, directed anteriorly, slightly projecting submedians; submedians triangular, directed posterolaterally, separated from medians by V-shaped notch; laterals narrowly triangular, directed anteriorly, separated from submedians by deep U-shaped notch, tip not reaching tip of submedians; inner orbital teeth low, triangular, directed posterolaterally. Anterolateral margins bearing 6 triangular teeth. Posterolateral margin sinuous, forming reentrant with posterior margin. Posterior margin nearly straight, about 34 percent carapace width. Vaulted metagastric and cardiac regions and medial parts of epibranchial regions ornamented with small granules irregularly arranged. Frontal, protogastric, mesogastric, epibranchial, and metagastric ridges present. A row of small granules sparsely arranged forming strongly convex arc on hepatic, anterolateral, and distal epibranchial regions.

Etymology: The trivial name honors Y. Kinugawa, who collected the type specimens and generously donated them to the Mizunami Fossil Museum.

Description: Carapace wider than long, length about 75 percent carapace width, widest at position of last anterolateral spine; carapace gently vaulted transversely and longitudinally. Front-orbital margin about 55 % carapace width. Median frontal teeth broadly triangular, directed anteriorly, slightly projecting submedian teeth. Submedian frontal teeth also triangular, directed posterolaterally, separated from median frontal teeth by V-shaped notch. Lateral frontal teeth narrowly triangular, directed anteriorly, separated from submedian frontal teeth by deep U-shaped notch, tip not reaching tip of submedian frontal teeth. Inner orbital teeth low, triangular, directed posterolaterally. Upper orbital margin concave with 2 closed fissures. Anterolateral margins bearing 6 triangular teeth including outer orbital tooth; 1st (=outer orbital tooth) and 2nd teeth directed anteriorly; third, fourth, and fifth teeth directed posterolaterally; sixth tooth directed laterally; third and fourth teeth larger than others. Posterolateral margin sinuous, forming reentrant with posterior margin. Posterior margin nearly straight, about 34 percent carapace width. Dorsal surface with poorly defined regions; anterior half nearly smooth, but posterior half, vaulted metagastric and cardiac regions and medial parts of epibranchial regions ornamented with small granules, which are irregularly arranged. Frontal ridges short, obtuse, medially interrupted. Protogastric and mesogastric ridges poorly defined, transverse, finely granulated, medially interrupted. Epibranchial ridges finely granulated, forming convex arc, not joining transverse, finely granulated metagastric ridge. A row of small granules sparsely arranged forming strongly convex arc on hepatic, anterolateral, and distal epibranchial regions.

Pereiopods poorly preserved. Ventral aspects unknown.

Remarks: The present new species resembles the extant species, *Charybdis* (*Charybdis*) *sagamiensis* Parisi, 1916, but differs in that the carapace is much wider than long and the metagastric and cardiac regions and medial parts of the epibranchial regions are ornamented with small, irregularly arranged granules. The unnamed fossil species of *Charybdis*, associated with carapaces, have been known from the Ananai Formation (Mimoto, ed., 1986) and Pliocene Shimajiri Group (Karasawa and Nobuhara, 2008); however, the present specimens cannot be compared with these specimens because of poorly preserved carapaces within latter specimens.



Fig. 2. *Charybdis* (*Charybdis*) *kinugawai* Karasawa, new species, MFM142021 (holotype), carapace, dorsal view. Scale bar = 1 cm.

References

- Berggren, W. A., F. J. Hilgen, C. G. Langerie, D. V. Kent, J. D. Obradovich, I. Raffi, M. E. Raymo, N. J. Shackleton (1995), Late Neogene chronology: New perspectives in high resolution stratigraphy. Geological Society of America, Bulletin, 107, 1272–1287.
- Cohen, K. M. and P. L. Gibbard (2011), Global chronostratigraphical correlation table for the last 2.7 million years, v. 2011. <http://www-qpg.geog.cam.ac.uk/>
- Dana, J. D. (1852), Macroura. Conspectus Crustaceorum &. Conspectus of the Crustacea of the Exploring Expedition under Capt C. Wilkes, U.S.N. Proceedings of the Akademy of Natural Sciences of Philadelphia, 6, 10–28.
- De Haan, W. (1833–1850), Crustacea. In Siebold, P. F. von, Fauna Japonica sive Descriptio Animalium, Quae in Itinere per Japoniam, Jussu et Auspicis Superiorum, qui Summum in India Batava Imperium Tenent, Suscepito, Annis 1823–1830 Collegit, Noitis, Observationibus et Adumbrationibus Illustravit. Leiden, Lugduni-Batavorum. i–xvii, i–xxxii, ix–xvi, 1–243, plates A–J, L–Q, 1–55.
- De Saint Laurent, M. (1979), Vers une nouvelle classification des Crustacés Décapodes Reptantia. Bulletin de l'Office National des Pêches République Tunisienne, Ministere de l'Agriculture, 3, 15–31.
- De Saint Laurent, M. (1980), Sur classification et phylogenie des Crustaces Décapodes brachyoures. I. Podotremata Guinot, 1977, et Eubrachyura sect. nov. Comptes Rendus Hebdomadaires des Seances de l'Academie des Sciences, series D, 290, 1265–1268.
- Hu, C.-H., and H.-J. Tao (2000), Crustacean fossils from southern Taiwan. Petroleum Geology of Taiwan, 34, 105–195.
- Imazumi, R. (1958), An interesting crustacean remain, *Ctenocheles sujakui* n. sp. from the Paleogene of Kyushu Japan. Transactions and Proceedings of the Palaeontological Society of Japan, N. S., 32, 299–304, pl. 44.
- Iwai, M., Y. Kondo, N. Kikuchi, and M. Oda (2006), Overview of stratigraphy and paleontology of Pliocene Tonohama Group, Kochi Prefecture, southwest Japan. Journal of the Geological Society of Japan, 112 (Supplement), 27–40.
- Karasawa, H. (1993), Cenozoic decapod Crustacea from southwest Japan. Bulletin of the Mizunami Fossil Museum, 20, 1–92.
- Karasawa, H. (1997), A monograph of Cenozoic stomatopod, decapod, isopod and amphipod Crustacea from west Japan. Monograph of the Mizunami Fossil Museum, 8, 81 p.
- Karasawa, H. and Y. Fudouji. (2000), Palaeogene decapod Crustacea from the Kishima and Okinoshima Groups, Kyushu, Japan. Paleontological Research, 4, 239–253.
- Karasawa, H. T. Nakagawa, and K. Kaede (2011), Axiidea and Brachyura (Decapoda) from the Miocene Shimo Formation, Uchiura Group, Fukui Prefecture, Japan. Bulletin of the Mizunami Fossil Museum, 37, 31–36.
- Karasawa, H. and T. Nobuhara (2008), Decapoda and Isopoda (Crustacea) from the Pliocene Shimajiri Group in the Miyako-jima island, Ryukyu, Japan. Bulletin of the Mizunami Fossil Museum, 34, 23–30.
- Karasawa, H., Schweitzer, C. E., and Feldmann, R. M. (2008), Revision of Portunoidea Rafinesque, 1815 (Decapoda: Brachyura) with emphasis on the fossil genera. Journal of Crustacean Biology, 28, 82–127.
- Katto, J. and K. Masuda (1993), Molluscan fossils from the Tonohama Group, famous for the legend of the Institute of the “Inedible Shells of Aki”. Sakawa Chisitukan Tenji Kaisetu Tokusyu, 1, 51 p.
- Kishinouye, K. (1926), Two rare and remarkable forms of macrurous Crustacea from Japan. Annotationes Zoologicae Japonenses, 11, 63–70.
- Latreille, P. A. (1802–1803), Histoire naturelle, générale et particulière, des crustacés et des insectes. Volume 3, 467 p. F. Dufart, Paris.
- Manning, R. B. and Felder, D. L. (1991), Revision of the American Callianassidae (Crustacea: Decapoda: Thalassinidea). Proceedings of the Biological Society of Washington, 104, 764–792.
- Matsubara, T. (2004), Catalogue of the Pliocene Mollusca from the Tonohama Group in Kōchi Prefecture, Shikoku, Japan, in the Museum of Nature and Human Activities, Hyogo (Takao Sendō Collection). Nature and Human Activities, 8, 49–95.
- Milne Edwards, H. (1834), Histoire naturelle des Crustacés, comprenant l'anatomie, la physiologie et la classification de ces animaux. Librairie Encyclopédique de Roret. Vol. 1. Paris. 468 p.
- Mimoto, K. (2001), *Macrocheira* sp. (Decapoda) and other crustacean fossils from the Pliocene Nobori Formation, Shikoku, Japan. Chigakukenkyū, 50, 131–135.
- Mimoto, K. (ed.) (1986), Fossils in Kochi Prefecture—illustrations of the specimens kept by members of the Kochi Fossil Society—. Kochi Fossil Society. 53 p.
- Ng, P. K. L., C.-H. Wang, P.-H. Ho, and H.-T. Shih (2001), An annotated checklist of brachyuran crabs from Taiwan (Crustacea: Decapoda). National Taiwan Museum Special Publication Series, 11, 86 p.
- Ortmann, A. (1893), Abtheilung: Brachyura (Brachyura genuina Boas), II. Unterabtheilung: Cancroidea, 2. Section: Cancerinea, 1. Gruppe: Cyclometopa. Die DecapodenKrebse des Strassburger Museums, mit besonderer Berücksichtigung der von Herrn Dr. Döderlein bei Japan und bei den Liu-Kiu-Inseln gesammelten und zur Zeit im Strassburger Museum aufbewahrten Formen, VII. Theil. Zoologische Jahrbücher, Abtheilung für Systematik, Geographie, und Biologie der Thiere, 7, 411–495.
- Parisi, B. (1916), Il Decapodi giapponesi del Museo di Milano. II. Cyclometopa, Atti della Società Italiana di Scienze Naturale di Milano, 55, 153–190.
- Paul'son, O. M. (1875 [reprint 1961]), Studies on Crustacea of the Red Sea with notes regarding other seas. Part I. Podophthalmata and Edriophthalmata (Cumacea). 164 p., 21 pls. The Israel Program for Scientific Translations, Jerusalem.
- Rafinesque, C. S. (1815), Analyse de la nature, ou tableau de l'universe et des corps organisés. 224 p. L'Imprimérie de Jean Barravecchia, Palermo, Italy.
- Sakai, K. (2010), Axiidea of the world and a reconsideration of the Callianassoidea (Decapoda, Thalassinidea, Callianassida). Crustaceana Monographs, 13, i–x +616 p.
- Schweitzer, C. E. and Feldmann, R. M. (2000), Reevaluation of the Cancridae Latreille, 1803 (Decapoda: Brachyura) including three new genera and three new species. Contributions to Zoology, 69, 223–250.