

# Recent and fossil Clinocardiinae (Bivalvia, Cardiidae) of the World. V. Genus *Ciliatocardium* Kafanov, 1974 (part 1)

Alexander I. Kafanov

Institute of Marine Biology, Far East Branch of the Russian Academy of Sciences, 17  
Palchevsky Str., Vladivostok 690041, Russia <Kafanov@mail.primorye.ru>

## Abstract

A taxonomic review of the genus *Ciliatocardium* Kafanov, 1974. In the modern fauna it is composed by *C. ciliatum ciliatum* (Fabricius, 1780), *C. ciliatum nordenskioldi* Kafanov, 1981, *C. ciliatum ochotense* Kafanov, 1981, *C. ciliatum pubescens* (Couthouy, 1838), *C. ciliatum tchuktchense* Kafanov in Scarlato, 1981 and *C. likharevi* Kafanov in Scarlato, 1981. Pleistocene *C. ciliatum dawsoni* (Stimpson, 1863) is known from Arctic coast of North America. *C. ciliatum* has also two chronosubspecies: Miocene *C. ciliatum maruyamense* Kafanov et Savizky, 1982 and Pliocene *C. ciliatum chikagawaense* (Kotaka, 1950). Key for identification of all Recent and fossil species and subspecies is given.

*Key words:* Bivalvia, Cardiidae, Clinocardiinae, *Ciliatocardium*, Recent, Cenozoic, Arctic Ocean, North Pacific, North Atlantic

## Genus *Ciliatocardium* Kafanov, 1974

*Ciliatocardium* Kafanov, 1974: 1469; Kafanov, 1980: 310.

*Type-species (OD)* : *Cardium ciliatum* Fabricius, 1780; Greenland, Recent.

Shell medium- and small sized or minute, oval-rounded or truncated-trigonal. Radial ribs about 17-52 in number, often widely arranged on the anterior area of external valve surface. Ribs triangular or tectate (roof-shaped) in cross-section. Crests of ribs with longitudinal rows of thin ciliated periostracal fringes (in adult and poorly preserved fossil shells ribs may be differently smoothed). Small spiniform lobes observed sometimes in juveniles on crests of ribs; sometimes these lobes are available and for the adult shells. The tooth *AI* without longitudinal ridge on its dorsal surface. Lunula oblong and cordiform or absent. Escutcheon if present narrow and lanceolate. Distal part of foot with narrow ventral sulculus surrounded on both sides by longitudinal rows of delicate papillae. Labial palps long but less than a half of the length of the inner demibranch.

Paleocene - Recent; cold and temperate waters of the northern hemisphere: northwestern (south to Kyushu, Japan) and northeastern (south to California, U.S.A.) Pacific; Arctic Ocean and North Atlantic (south to Cape Cod, Massachusetts, U.S.A., Iceland and southern Norway; in Pliocene to England).

## Key to the species and subspecies

1. Average ribs number more than 35.....2.  
Average ribs number less than 35.....13.
2. Average H/L ratio 0.95 or more.....3.  
Average H/L ratio less than 0.95.....6.
3. Umbo medium wide; ribs with rather straight  
shape of slope.....4.  
Umbo wide; ribs rather convex-shaped..... 5.
4. Average B/L ratio about 0.29; ribs 28-35 in number  
....*C. ciliatum maruyamense* Kafanov et Savizky, 1982.  
Average B/L ratio about 0.36; ribs 30-40 in number  
.....*C. ciliatum chikagawaense* (Kotaka, 1950).
5. Average A/L ratio about 0.48; average B/L ratio about

- 0.38; ribs 44-49 in number..... *C. coosense* (Dall, 1909).  
Average A/L ratio about 0.41; average B/L ratio about  
0.27; ribs 33-45 in number..... *C. smekhovi* Kafanov et  
Savizky, 1982.
6. Average ribs number more than 40.....7.  
Average ribs number less than 40.....8.
7. Ribs 42-48 in number; average H/L ratio about 0.76  
... *C. ermanense* Sinelnikova in Sinelnikova *et al.*, 1976  
Ribs 45-52 in number; average H/L ratio about 0.88  
..... *C. carlsbergianum* Kafanov, sp. nov.
8. Average H/L ratio less than 0.90.....9.  
Average H/L ratio more than 0.90.....10.
9. Average H/L and A/L ratios 0.88 and 0.32 accordingly;  
shell moderately oblique and faintly bent  
..... *C. brooksi* (Clark, 1932).  
Average H/L and A/L ratios 0.80 and 0.43 accordingly;  
shell non-oblique and unbent  
..... *C. zhidkova*e Kafanov in Arkhipova *et al.*, [1994].
10. Adult shell minute or small-sized.....11.  
Adult shell medium-sized.....12.
11. Average B/L and A/L ratios 0.23 and 0.47 accordingly;  
ribs 36-44..... *C. ainuanum* (Yokoyama, 1927).  
Average B/L and A/L ratios 0.29 and 0.41 accordingly;  
ribs 35-42..... *C. asagaiense* (Makiyama, 1934).
12. Shell equilateral, non-oblique and unbent; anterior  
branch of dorsal margin concave; average nymph  
length is no more than 1/5 of the shell length  
..... *C. likharevi* Kafanov in Scarlato, 1981.  
Shell at least faintly oblique and bent, rather subequi-  
lateral; anterior branch of dorsal margin straight or  
faintly convex; average nymph length is more than  
1/5 of the shell length  
..... *C. ciliatum* (Fabricius, 1780) [Recent subspecies].
13. Ribs 27 or less.....14.  
Ribs more than 27.....20.
14. Adult shell minute.....15.  
Adult shell small- or medium-sized.....16.
15. Ribs 17-20 in number; average H/L ratio about 0.93  
..... *C. kirkerense* (Clark, 1918).  
Ribs 21-25 in number; average H/L ratio about 0.90  
..... *C. reedi* (Loel et Corey, 1932).
16. Average H/L ratio more than 0.90.....17.  
Average H/L ratio less than 0.90.....18.
17. Average B/L and A/L ratios 0.27 and 0.46 accordingly;  
ribs 22-25..... *C. matchgarensis* (Makiyama, 1934).  
Average B/L and A/L ratios 0.34 and 0.42 accordingly;  
ribs 20-26..... *C. yamasakii* (Makiyama, 1934).
18. Average B/L ratio less than 0.25  
..... *C. mutuense* (Nomura et Hatai, 1936).  
Average B/L ratio more than 0.25.....19.
19. Average B/L and A/L ratios 0.36 and 0.46 accordingly  
..... *C. sachalinense* (Khramova, 1962).  
Average B/L and A/L ratios 0.31 and 0.43 accordingly  
..... *C. uyemurai* (Kanehara, 1937).
20. Ribs less than 30 in number.....21.  
Ribs 30 and more in number.....24.
21. Average H/L ratio more than 1.10  
..... *C. moorae* Kafanov, sp. nov.  
Average H/L ratio less than 1.10.....22.
22. Average B/L ratio more than 0.30  
..... *C. hataii* (Hayasaka, 1956).  
Average B/L ratio less than 0.30.....23.
23. Average A/L ratio about 0.45  
..... *C. makiyamae* (Kamada, 1962).  
Average A/L ratio about 0.49  
..... *C. tigilense* (Slodkewitsch, 1938).
24. Average H/L ratio more than 0.95.....25.  
Average H/L ratio less than 0.95.....26.
25. Ribs 33-37 in number  
..... *C. shinjiense* (Yokoyama, 1923).  
Ribs 24-32 in number  
..... *C. yakatagense* (Clark, 1932).
26. Average H/L ratio more than 0.86.....27.  
Average H/L ratio less than 0.86.....28.
27. Adult shell medium-sized; umbo wide  
..... *C. iwataense* (Chinzei, 1959).  
Adult shell small-sized; umbo narrow  
..... *C. obstinatus* (Barinov in Gladenkov *et al.*, 1992).
28. Average H/L, B/L and A/L ratios 0.85, 0.25 and 0.42  
accordingly; ribs 27-36 in number  
..... *C. evssevi* Kafanov et Savizky, 1982.  
Average H/L, B/L and A/L ratios 0.82, 0.29 and 0.44  
accordingly; ribs 25-33 in number  
..... *C. schmidtii* (Khramova, 1962).

***Clitocardium ciliatum* (Fabricius, 1780)<sup>1</sup>**  
(Pls. 1-6; pl. 7, figs. 1-8)

*English:* Iceland cockle, hairy cockle; *Japanese:* Kokera-  
ishikage-gai; *Icelandian:* báruskel; *local in Greenland:*

kirksoarsak.

*Cardium ciliatum* Fabricius, 1780: 410.

*Cardium californiense* var. *comoxense* Dall, 1900: 1093.

*Laevicardium decoratum* (Grewingk): Grant and Gale, 1931, pl.  
19, fig. 12.

<sup>1</sup> Synonymy and bibliography herein is given for fossil specimens mainly, for Recent ones see distinct subspecies.

Table 1. Two-variable matrix for paired Comparison: (Student's t-test) of the power function coefficients **b** in four subspecies of *Ciliatocardium ciliatum*.

Variables	Subspecies	II	III	IV
H, L	I	4.27*** (69; 63)	3.10** (69; 69)	1.48 (69; 16)
A, L	I	2.26* (69; 63)	1.53 (69; 69)	1.02 (69; 16)
B, L	I	6.16*** (69; 63)	1.87 (69; 69)	0.96 (69; 16)
n, L	I	2.71** (69; 63)	-	0.42 (69; 16)
H, L	II	-	0.24 (69; 63)	0.29 (63; 16)
A, L	II	-	3.44*** (69; 63)	0.36 (63; 16)
B, L	II	-	3.00** (69; 63)	1.76 (63; 16)
n, L	II	-	-	1.56 (63; 16)
H, L	III	-	-	0.10 (69; 16)
A, L	III	-	-	1.50 (69; 16)
B, L	III	-	-	0.07 (69; 16)
n, L	III	-	-	-

Power function is  $X = aL^b$ , where **X** are valve height (**H**), length of the valve anterior part (**A**), valve convexity (**B**) and length of nymph (**n**) accordingly. Significance: \*P(0.1), \*\* P(0.01), \*\*\*P(0.001). Subspecies: I - *C. ciliatum ciliatum*; II - *C. ciliatum nordenskiöldi*; III - *C. ciliatum tchuktchense*; IV - *C. ciliatum ochotense*. Degree of freedoms for compared sets are given in brackets. *C. ciliatum pubescens* is not included in this table because its morphometric characters significantly differ from the parameters of normal (Gaussian) distribution.

*Papyridea sertunayana* Kogan (in litt.): Slodkewitsch, 1938: 405, pl. 82, figs. 3, 3a.

*Clinocardium comoxense* Dall, 1900: Keen, 1954: 19, pl. 1, figs. 5, 7, 8.

*Laevicardium* cf. *decoratum* "(Grewingk, 1850)": Ilyina, 1960, pl. 2, fig. 1.

*Clinocardium decoratum* "Grewingk, 1850": Zhidkova *et al.*, 1968, pl. 20, fig. 16.

*Clinocardium sachalinense* Khranova, 1962: Zhidkova *et al.*, 1972, pl. 11, fig. 1.

?*Profulvia sertunayana* Slodkewitsch, 1938: Kafanov, 1980: 316.

*Clinocardium ciliatum padimeicum* Merklin and Zarkhidze in Merklin *et al.*, 1979: 44, pl. 7, figs. 10, 11.

*Ciliatocardium olchovensis* Petrov, 1982: 104, pl. 19, figs. 1, 2, 4, 7, 8.

*Ciliatocardium ciliatum* (Fabricius): Gladenkov *et al.*, 1980: 43, pl. 8, figs. 1-3; Kafanov, 1980: 312; Kafanov and Savizky, 1982: 55; Petrov, 1982: 103, pl. 19, figs. 3, 5, 6.

?*Ciliatocardium ciliatum* (Fabricius): Gladenkov *et al.*, 1992, pl. 12, fig. 2.

**Bibliography:** Chinzei, 1959; Clarke *et al.*, 1972; Dall, 1919, 1924a, b; Durham and MacNeil, 1967; Gladenkov, 1972; Grant and Gale, 1931; Grigorenko *et al.*, 1967; Hopkins, 1959; Hopkins *et al.*, 1965; Iwai and Siobara, 1969; Kafanov, 1981a; Kotaka and Noda, 1967; Kuroda, 1931; Marinovich, 1983; Marinovich and Powell, 1989; Merklin *et al.*, 1962; Noda, 1973; Petersen and Kristensen, 1991; Richards, 1962; Schlesch, 1924; Soot-Ryen, 1932; Uozumi, 1962; Zinbo, 1973.

**Diagnosis:** Shell medium-sized, trigonal-subcircular or trigonal-suborbicular, high (average H/L ratio about 0.87-

0.99), slightly subequilateral (average A/L ratio about 0.40-0.43), medium convex (average B/L ratio about 0.29-0.32), faintly oblique and faintly bent, rather thick-walled. Keel twist not present. Ribs 26-44 (34-38 in average) in number, triangular-shaped, with straight or concave shape of slope. Quite often hinge is strongly weakened at the expense of a reduction of a tooth AIII and cardinals. Usually length of nymph is more than 1/5 of the shell length.

**Comparison:** Recent sibling species, *C. likharevi* Kafanov in Scarlato, 1981, differs by concave anterior branch of hinge margin, by non-oblique and unbent shell and by relatively shorter nymph (in average no more than 1/5 of the shell length).

**Taxonomical notes:** Pleistocene *Cardium californiense* var. *comoxense* Dall, 1900 (pl. 1, figs. 13-16), *Clinocardium ciliatum padimeicum* Merklin et Zarkhidze in Merklin *et al.*, 1979 and *Ciliatocardium olchovensis* Petrov, 1982 (pl. 1, fig. 17) by nothing differ from Recent *Ciliatocardium ciliatum* and should be considered as synonyms of the last. Possible synonym also is *Papyridea sertunayanum* Slodkewitsch, 1938 (pl. 4, fig. 8) described on the deformed shell.

Dall (1900) marks extremely wide variability of *C. ciliatum* and does not consider possible to establish any discrete groups inside this species. However, elementary statistical analysis of main conchometric parameters (Table 1) permits separate five groups, to which quite approach criteria of subspecies rank taxa: nominative subspecies, *C. ciliatum nordenskiöldi* Kafanov, 1981, *C. ciliatum ochotense* Kafanov, 1981, *C. ciliatum pubescens* (Couthouy, 1838) and *C. ciliatum tchuktchense* Kafanov in Scarlato, 1981. Pleistocene *C. ciliatum dawsoni* (Stimpson, 1863) can be considered also as separate subspecies and others two chronosubspecies are submitted by Pliocene *C. ciliatum chikagawaense* (Kotaka, 1950) and Miocene *C. ciliatum maruyamense* Kafanov et Savizky, 1982. Following key may be used for determination of Recent and Pleistocene subspecies:

1. Average H/L ratio about 0.88-0.95; umbo moderately high and wide; if external valve surface is corroded that is usual on umbonal area only.....2.
2. Average H/L ratio about 0.93-0.96; umbo low and wide (if high that tapering); external valve surface is strong corroded.....4.
2. Posteroventral corner is not expressed; shell rather equilateral

- .....*C. ciliatum pubescens* (Couthouy, 1838).  
Posteroventral corner expressed in a different degree;  
shell subequilateral.....3.
3. Slightly curved posterior branch of hinge margin  
smoothly passes in posterior margin of valve; average  
A/L ratio about 0.41; average B/L ratio about 0.31  
.....*C. ciliatum ciliatum* (Fabricius, 1780).  
Almost straight posterior branch of hinge margin pass-  
es in posterior margin of valve with an appreciable  
corner; average A/L ratio about 0.43; average B/L  
ratio about 0.30  
....*C. ciliatum tchuktchense* Kafanov in Scarlato, 1981.
4. Posteroventral corner well expressed, slightly pointed  
and extended backward; umbo high and narrow, well  
marked by anterior umbonal cavity; average B/L  
ratio about 0.22  
.....*C. ciliatum dawsoni* (Stimpson, 1863).  
Posteroventral corner rounded and extended down-  
ward; umbo low and wide, not clearly marked by an-  
terior umbonal cavity; average B/L ratio more than  
0.22.....5.
5. Ribs flattened, wide; ciliated periostracal fringes on  
the ribs are poorly advanced and on the medial area  
almost always are absent  
.....*C. ciliatum ochotense* Kafanov, 1981.  
Ribs rather high, narrow, pointed, concave-shaped; cil-  
iated periostracal fringes on the ribs are well  
advanced and also on the medial area  
.....*C. ciliatum nordenskiöldi* Kafanov, 1981.

*Geographical distribution* (Fig. 1): One of the most  
usual Bivalvia species in cold (Arctic Ocean) and temper-

ate waters of the northern hemisphere. Southern limits of  
distribution: North Atlantic - east Finnmarken, Norway,  
southwest coast of Färöer Islands, Denmark, south Ice-  
land, south Greenland and Cape Cod, Massachusetts,  
U.S.A.; North Pacific - East Korea Bay, Tajima, Hyogo  
Prefecture, Japan, Otsuchi Bay, northeastern Honshu,  
Japan, Commander and Aleutian Islands, Puget Sound,  
Washington, U.S.A.

*Geochronological distribution*: The earliest fossil occur-  
rences of *C. ciliatum* are lower Middle Miocene Ulegorsk  
Horizon (Aleksandrovskaia and/or Sertunayskaia Suite)  
of west Sakhalin, Russia. In West Kamchatka it is known  
not earlier than middle Miocene Etolonskaia Suite  
(Grigorenko *et al.*, 1967). It is reported also from Miocene  
Ogawara Formation of Aomori Prefecture, Honshu.  
*Ciliatocardium ciliatum maruyamense* Kafanov et  
Savizky, 1982 is distributed in upper Miocene deposits of  
Sakhalin and *C. ciliatum chikagawaense* (Kotaka, 1950) is  
the usual representative of Pliocene Omma fauna in  
Japan. In Eastern Kamchatka *C. ciliatum* is known from  
Pliocene Ust'-Limimteveyamskaia Suite of Karaginsky  
Island. Kanehara's (1937) data for Oligocene Iwaki  
Formation of Northeast Honshu are erroneous.

Analogous data for Pacific coast of North America  
belongs only for Pliocene Beringian strata of Pribilof  
Islands, Alaska, U.S.A. (Grant and Gale, 1931) though  
similar forms are known from upper Miocene Tachilni  
Formation of southwest Alaska (Marincovich, 1983). It is  
common also in Pliocene and Pleistocene deposits of  
Chukotka Peninsula, Arctic coast of Russia, Alaska,  
Iceland (Tjörnes Crag) and England (Icenian).




Fig 1. Geographical distribution of Recent *Ciliatocardium ciliatum* (Fabricius, 1780). Subspecies: 1 - *C. ciliatum ciliatum* (Fabricius, 1780), 2 - *C. ciliatum pubescens* (Couthouy, 1838), 3 - *C. ciliatum nordenskiöldi* Kafanov, 1981, 4 - *C. ciliatum tchuktchense* Kafanov in Scarlato, 1981, 5 - *C. ciliatum ochotense* Kafanov, 1981, 6 - specimens from the intergradation zone between subspecies.

***Ciliatocardium ciliatum ciliatum***  
(Fabricius, 1780)

(Pl. 1, figs. 1-11, 18; pl. 2, figs. 1-13)

- Cardium ciliatum* Fabricius, 1780: 410; Römer, 1869: 43 (part.), Taf. 7, Fig. 5, 6 non Taf. 10, Fig. 1-3; Mörch, 1870: 27 (part.); Leche, 1878:21 (part.); Sars, 1878: 46, pl. 5, figs. 4a, b; Herzenstein, 1885: 652; Posselt, 1895: 70 (part.?); Posselt and Jensen, 1898: 57 (part.?); Friele and Grieg, 1901: 33 (part.); Knipowitsch, 1902: 93 (part.); Hägg, 1905: 51 (part.); Odhner, 1910: 18; Jensen, 1912: 79, pl. 3, fig. 10 (juv.); Odhner, 1915: 118 (part.); Grieg, 1924: 7 (part.); Johnson, 1926: 24 (part.); Thiele, 1928: 577, 621 (part.); Pirozhnikov, 1937: 83, pl. 38, fig. 1; Madsen, 1949: 63; Thorson, 1951: 80; Óskarsson, 1952: 78, text-fig. 69 (reproduced from: Sars, 1878, pl. 5, fig. 4); Ockelmann, 1958: 118 (part.?); Petersen, 1978, text-fig. 1C.
- Cardium islandicum* Chemnitz, 1782: 200, pl. 19, figs. 195, 196, nonbinom. [Official index..., 1958: 5, Direction 1].
- Cardium islandicum* Bruguière, 1789: 222; Gmelin, 1791: 3252; Hancock, 1846: 336 (part.?); Middendorff, 1849: 38 (part.?); Packard, 1867: 278; Smith, 1877: 141 (part.?); Kobelt, 1888: 367 (part.); Rogers, 1920: 358 (part.).
- Cardium edule*, non Linnaeus, 1758: Mohr, 1786: 128; Jeffreys, 1864: 287.
- Cardium icelandicum* "Gmelin": Wood, 1825: 26, pl. 5, fig. 27; Sowerby, 1841b: 2, no. 19; Reeve, 1845: sp. 54, pl. 11.
- ?*Cardium boreale* Broderip et Sowerby, 1829: 368 non Reeve, 1845: sp. 131, pl. 22.
- ?*Cardium arcticum* Sowerby, 1841a: 106; Sowerby, 1841b: 2, no. 27, fig. 26.
- Cardium (Serripes) islandicum* Bruguière: Adams and Adams, 1858: 456.
- Cardium hayesii* Stimpson, 1864: 142; Packard, 1867: 278, pl. 7, fig. 14.
- Cardium (Cerastoderma) ciliatum* Fabricius: Adams and Adams, 1858: 456; Dall, 1900: 1096 (part.); Dall, 1901: 386 (part.); Whiteaves, 1901: 128 (part.); McLean, 1939: 162, pl. 24, figs. 1, 2; Filatova, 1948: 432 (part. non pl. 109, fig. 1).
- Cardium (Cerastoderma) ciliatum* Bruguière: Dautzenberg and Fischer, 1912: 448 (part.; comprehensive synonymy until 1902).
- Clinocardium ciliatum* (Fabricius): Keen, 1936: 120 (part.); LaRocque, 1953: 63 (part.); Keen, 1954: 20 (part.); Soot-Ryen, 1958: 22 (part.?); Bousfield, 1960: 28 (part.), pl. 6, fig. 70; Allen, 1965: 938 (part.); Abbott, 1968: 228, text-fig. 6?; Nordsieck, 1969: 98, pl. 15, fig. 54.30; Voskuil, 1989: 273, fig. 24.
- Corculum (Keenocardium) ciliatum* (Fabricius): Fischer-Piette, 1977: 134 (part.).
- Ciliatocardium ciliatum ciliatum* (Fabricius): Kafanov, 1981a: 49, figs. 2a, b.
- Ciliatocardium ciliatum* (Fabricius): Golikov and Scarlato, 1977: 368 (part.); Naumov *et al.*, 1987: 247, fig. 154.

*Bibliography:* Ecology, life cycle, morphology, distribu-

tion, fisheries: Andrews, 1972; Brøgger, 1901; Brotskaya and Zenkevich, 1939; Brotskaya *et al.*, 1963; Clarke, 1974; Dall, 1924b; Derjugin, 1915, 1928; Fedyakov, 1986; Filatova, 1957; Gorbunov, 1952; Jensen, 1901; Kaufman, 1974, 1977; Kuznetsov, 1963, 1970; Kuznetsov, 1960, 1961, 1963; Messjatzaw, 1930, 1931; Miloslavskaya, 1958a, b; Petersen, 1968, 1978; Powles, 1965; Spärck, 1933, 1936, 1937; Thorson, 1933, 1934, 1936, 1957; Tseeb, 1964; Tsikhon-Lukanina, 1987; Turpaeva, 1948, 1953; Vibe, 1939, 1950; Zatsepin, 1962; Zatsepin and Filatova, 1961; Zenkevich, 1947, 1951, 1963; Zenkevich and Brotskaya, 1937; Zhirmunsky, 1969.

*Etymology:* The name is derived from the new Latin *cilia*, eyelashes.

*Type-materials:* *Cardium ciliatum* Fabricius, 1780. Lectotype (designated by Kafanov, 1980: 312) and paralectotype are stored in ZMK (reg. no. absent). Type locality: Greenland (most possibly, southwest). Lectotype dimensions: L = 51.8; H = 47.6; B = 17.1; A = 21.0.

*Cardium islandicum* Bruguière, 1789 is based on the nonbinominal name of Chemnitz (1782). Type locality: Iceland. Dall (1900) marks that many old authors attributed authorship of this species to Linnaeus and quite often even with the link to 12th edition of "Systema Naturae", where such name is not mentioned. Tryon's (1873: 1975) indication on "*Cardium islandicum* Linnaeus, edit. 12, p. 1124 "concerns to *Cardium pectinatum* Linnaeus, 1758.

*Cardium arcticum* Sowerby, 1841. Possible syntype - BMNH no. 1975618 (Pl. 1, figs. 9-11; pl. 4, fig. 10). Type locality: "Arctic seas".

*Cardium hayesii* Stimpson, 1864. Type materials are lost; Dance (1966) reports that Stimpson's collection was destroyed at the Chicago fire in 1871. Type locality: Disko Island, west Greenland.

*Description* (see also Table 2): Shell medium-sized (maximum to 65 mm), trigonal-subcircular or trigonal-suborbicular, high (average H/L ratio about 0.95), slightly subequilateral (average A/L ratio about 0.41), medium convex (average B/L ratio about 0.31), faintly oblique but unbent, rather thick-walled. Posterior branch of hinge margin smoothly passes in posterior margin of valve; posteroventral corner is poorly expressed and usually not expressed down- or backward. Growth rings well marked. Umbo marked mainly by anterior umbonal cavity, wide and medium height, often slightly corroded. Ribs 32-38 (34-36 in average) in number, rather low and wide. On the medial

area ribs are not bent, regularly-triangular in cross-section, separated by narrower intercostal spaces. Become closer to anterior area ribs increase height and intercostal spaces extend. The most forward four ribs are pleated bent, widely placed and are shared by 1.5-2 times wider intercostal spaces. Last two ribs on the escutcheon area have character of not clear wrinkled folds. Ciliated periostracal fringes on the ribs are poorly advanced. As a rule, the hinge is not reduced. Lunula lengthened, slightly pressed.

*Comparison:* The most similar *C. ciliatum tchuktchense* Kafanov in Scarlato, 1981 differs by following characters: almost straight posterior branch of hinge margin passes in posterior margin of valve with an appreciable corner; average A/L ratio about 0.43 (versus 0.41 in *C. ciliatum ciliatum*); average B/L ratio about 0.30 (versus 0.31).

*Taxonomic notes:* The majority of the authors considers *Cardium boreale* Broderip et Sowerby, 1829 non Reeve, 1845 as a synonym of *Cardium ciliatum* Fabricius. However, because the original description by Broderip and Sowerby (1829) is very brief and ambiguous, and the type specimen, never illustrated, has been lost the status of this species remains doubtful (see: Nomina dubia).

*Ecology:* Commonly inhabits lower intertidal and subtidal zones, from 0 up to 700 m preferably on silts, sandy

Table 2. Statistical values of basic morphometric characters in Recent *Ciliatocardium ciliatum ciliatum* (Fabricius, 1780).

N = 71;  $\bar{y} = L = 33.0 \pm 1.9$ ;  $\bar{y}_y = 16.3 \pm 1.4$ ;  $As_y = + 0.001$ ;  $E_y = - 0.614$ ;

Parameters	H	A	B	n
$\bar{x}$	31.5±1.9	13.2±0.7	10.4±0.6	8.9±0.7
$\sigma_x$	15.9±1.3	6.1±0.5	5.3±0.4	5.7±0.5
$As_x$	+0.031	+0.013	+0.188	+0.272
$E_x$	-0.487	-0.499	-0.568	-0.439
$r$	0.998±0.000***	0.995±0.001***	0.992±0.002***	0.984±0.004***
$r_{ln}$	0.999±0.000***	0.997±0.001***	0.996±0.001***	0.993±0.002***
$a$	1.190	1.988	3.889	7.720
$b$	0.976±0.005	1.072±0.010	0.969±0.010	0.730±0.010
$x/y$	0.95±0.00	0.41±0.00	0.31±0.00	0.24±0.01
$\sigma_{x/y}$	0.03±0.00	0.02±0.00	0.02±0.00	0.05±0.00
$As_{x/y}$	+0.204	-0.202	+0.256	+0.228
$E_{x/y}$	-0.237	-0.637	-0.415	-0.263

Notation admitted here and in tables 3-7 is as follows: N - sampling volume;  $\mathbf{y(L)}$  and  $\mathbf{x}$  (H, A, B, n) - values of the argument and functions (mean ± SD);  $\mathbf{As}$  and  $\mathbf{E}$  - coefficients of asymmetry and excess; for significant differences of  $\mathbf{As}$  and  $\mathbf{E}$  from the parameters of normal (Gaussian) distribution and for significance of  $\mathbf{r}$  and  $\mathbf{r}_{ln}$ : \*P(0.1), \*\* P(0.01), \*\*\*P(0.001;  $\mathbf{r}$  - paired Pearson's correlation coefficient between the values of  $\mathbf{x}$  and  $\mathbf{y}$ ;  $\mathbf{r}_{ln}$  - paired Pearson's correlation coefficient between the logarithms of  $\mathbf{x}$  and  $\mathbf{y}$ ;  $\mathbf{a}$  and  $\mathbf{b}$  - constants of the power functions  $\ln y = \ln a + b \cdot \ln x$ ; quotation of the type  $\mathbf{x/y} = 0.89 \pm 0.00$  denotes that SD(0.005).

silts and sands but often occurs on mixed soils and on pebbles. Most frequently distributes on depth 30-150 m at near-bottom temperature + 1-3 during August and September. In European sub-Arctic it frequently inhabits the extremely large depths (Soot-Ryen, 1932) that most likely is caused by transport activity of sea ice (Fig. 2). It prefers salinity 33-35‰ though in White Sea maintains considerably lowered salinity (up to 24‰) and abnormal high near-bottom temperature (up to 16 in Onega Bay).

Near Iceland it is a mass species in *Yoldia hyperborea* community (45-162 m on soft grounds). In fjords *C. ciliatum* includes here in *Macoma calcarea* community (2-50 m on soft grounds). At southern Iceland is rather rare and inhabits fjords mainly. At east Greenland it is common in *Macoma calcarea* community where reaches average biomass 32.4 g/m<sup>2</sup> (maximum up to 134 g/m<sup>2</sup>). At Murmansk Coast of the Barents Sea prevails in *C. ciliatum - Arctica islandica* community (10-60 m on muddy sands) and is characteristic species in *Strongylocentrotus droebachiensis - Astarte crenata - Nicomache lumbricalis* community. In structure of main bottom communities of Barents Sea (*Nuculana pernula - Astarte borealis*; *C. ciliatum - Astarte*




Fig 2. The maximal depths of a finding of empty valves of *Ciliatocardium ciliatum* (Fabricius, 1780) in Greenland Sea and adjacent waters.

The area of thawing of ice in places of meeting of warm and cold currents (schematically according Lisitsyn, 1961) is shaded. Depths (in m): 1 - 2754; 2 - 1447; 3 - 3145; 4 - 2968; 5 - 2224; 6 - 2320; 7 - 2886; 8 - 2715.

*borealis*, *Portlandia arctica*, etc.) it prevails on biomass (in average 38.5 g/m<sup>2</sup>) and density (in average 8 specimens/m<sup>2</sup>) on depth 0-100 m. In White Sea is usual in internal sites of inlets (6-22 m on silts) in *Nuculana pernula* - *Arctica islandica* - *C. ciliatum* - *Astarte elliptica* community complex.

One of the main components in a diet of walruses, whales and of many benthophagous fishes: halibuts, flounders, haddock, wolffishes and other. It is quite edible for the man (on taste reminds *Spisula sachalinensis*) but is not used.

*Distribution* (Fig. 1): Southwestern (to north up to the Polar circle) coast and southeastern extremity of Greenland, northeastern coast of Newfoundland Island, Canada, Iceland, southwest coast of Färöer Islands, Denmark, north Norway (to south up to east Finnmarken), southwest Spitzbergen, southwest part of Barents Sea (from the southern extremity of West Spitzbergen up to Svyatoy Nos Cape, Kola Peninsula, Russia).

Wide zone of intergradation with *C. ciliatum nordenskiöldi* occupies central and southeastern parts of Barents Sea, southeastern and northwestern coasts of West Spitzbergen and Baydaratskaya Inlet of Kara Sea. On the American coast northern border of distribution of *C. ciliatum ciliatum* is not known precisely, but at northeastern Labrador, Canada, specimens of a transitive type are distributed.

*Materials*: 254 lots (more than 950 specimens): ZIsp, CAS, LACMNH.

### ***Ciliatocardium ciliatum dawsoni***

(Stimpson, 1863)

(Pl. 1, fig. 12; pl. 4, figs. 6, 7)

*Cardium dawsoni* Stimpson, 1863: 58, text-fig.

*Ciliatocardium ciliatum dawsoni* (Stimpson, 1863): Kafanov, 1981a: 51, fig. 2B.

*Etymology*: The name is derived in the honour of Sir John William Dawson (1820-1899), Canadian geologist who made numerous contributions to paleobotany and extended the knowledge of Canadian geology.

*Type-material*: It follows from original description that the type-material was stored at Museum of Smithsonian Institution [USNM] however it is absent nowadays here (J. Rosewater, pers. comm., Nov. 17, 1973). Besides Dance (1966) reports that Stimpson's collection was destroyed at the Chicago fire in 1871. Type locality: "on the beach at Cape Hope, on the southeast side of Hudson's Bay, having

in all probability been washed out of a Pleistocene deposit". Original figure dimensions: L = 41.5; H = 38.8; B = 12.7.

*Description* (from: Stimpson, 1863: 58-59): "Shell subovate, oblique, very inequilateral, and somewhat angular posteriorly; beaks small and much elevated; hinge thin; teeth weak, especially the posterior ones; ribs about thirty-five in number, in the anterior part of the shell narrower than their interspaces, in the middle and posterior parts broader and more flattened; ventral margin crenulated".

*Comparison*: This form appreciably differs from others subspecies of *C. ciliatum* by shell-shape markedly angular posteriorly (see Key for subspecies).

*Taxonomic notes*: I had two left valves received from CAS and labelled: "CAS no. 34343. *Clinocardium* cf. *californiense* (Deshayes), Arctic coast of Alaska, 71° 31' 20" N 156° 23' 00" W, about 169 m, G. D. Hanna, Aug. 14 1954" and "CAS no. 35052. Deadman Island, off Point Barrow, Sh. Sovalik, M. Brewer and G. D. Hanna, June 21 1957". These shells quite correspond to the original description of *Cardium dawsoni* and its image. Taking into account essential morphological differences of this form *C. ciliatum dawsoni* (Stimpson, 1863) can be considered as separate subspecies characteristic for Pleistocene deposits of Arctic coast of North America.

*Materials*: 2 lots (2 specimens): ZIsp.

### ***Ciliatocardium ciliatum nordenskiöldi***

Kafanov, 1981

(Pl. 3, figs. 1-4; pl. 5, figs. 1-13)

*Cardium ciliatum* Fabricius: Mörch, 1870: 27 (part.); Leche, 1878: 21 (part.); Leche, 1883: 443 (part.); Pfeffer, 1886: 9; Friele and Grieg, 1901: 33 (part.); Knipowitsch, 1902: 93 (part.); Knipowitsch, 1903: 52 (part.?); Hägg, 1905: 51 (part.); Odhner, 1915: 118 (part.); Grieg, 1924: 7 (part.); Thiele, 1928: 621 (part.).

*Cardium islandicum* Bruguière: Kobelt, 1888: 367 (part.).

*Cardium (Cerastoderma) islandicum* Bruguière: Dautzenberg and Fischer, 1912: 448 (part.);

*Ciliatocardium ciliatum* (Fabricius): Golikov and Scarlato, 1977: 368 (part.); Naumov and Fedyaev, 1990: 403.

*Clinocardium ciliatum* (Fabricius): Lubinsky, 1980: 38 (part?), pl. 8, fig. 1.

*Ciliatocardium ciliatum nordenskiöldi* Kafanov, 1981a: 52, figs. 2D, Zh-I.

*Bibliography*: Ecology, life cycle, morphology, distribution, fisheries: Filatova, 1957; Gorbunov, 1946; Messjatzaw, 1931; Popova, 1948.

*Etymology*: The name is derived in the honour of Nils

Otto Gustaf Nordenskjöld (1869-1928), Swedish geographer and explorer whose expedition to the Arctic was distinguished by the volume of its scientific findings.

*Type-material:* Holotype (reg. no. 1/200156) and paratypes (reg. nos. 200157-200273) in ZIsp. Type locality: Laptev Sea, Komsomol'skoy Pravdy Islands, between Angreya Island, 76°50' N 110°40' E and Samuila Island, 77°25' N 106°45' E, depth 37 m, mud, near-bottom temperature -1.2 °C; leg. E. E. Arngold, r/v "Vaygach", August 13(26), 1913.

Holotype dimensions: L = 40.3; H = 38.1; B = 11.7; A = 16.1.

*Description* (see Table 3): Shell medium-sized (usually not more than 50mm in length), suborbicular, high (average H/L ratio about 0.96), subequilateral (average A/L ratio about 0.42), medium convex, faintly oblique and faintly bent, thin-walled, often corroded and even perforated especially at umbonal area. Keel twist absent. Growth rings well marked, numerous. Umbo low, wide, blunt, badly designated by umbonal cavities. Ribs 27-39 (32-34 in average) in number, thin, not enough bent, concave-shaped in cross-section, widely placed, wrinkled, bearing distinct longitudinal rows of periostracal fringes or of spike-form tubercles in juvenile specimens. Hinge plate narrow. Teeth, especial *AI*, *3a* and *4b*, are essentially weakened. In area of cardinal teeth additional semiinternal ligament is present. The tooth *PI* is almost equal to nymph length. Internal surface of valve Brilliant, with clear negative ribs. Extremely rare specimens from western part of East Siberian Sea have a small and convex shell perforated at umbonal area and rather large spike-form tubercles on the ribs.

*Comparison:* The most similar *C. ciliatum ochotense* Kafanov, 1981 differs by shell-outline, by wide and flattened ribs; ciliated periostracal fringes on the ribs are poorly advanced and on the medial area almost always are absent among this subspecies.

*Ecology:* The range of vertical distribution makes from 9 up to 293 m however most frequently inhabits soft bottom on depth 20-40 m. Subfossil valves, most probable transported by ice, were found in Laptev Sea (77°25' N 118°17' E) on depth 869 m. The typical specimens are found out only at constantly negative near-bottom temperatures: from -0.4 up to -1.7 °C in Kara Sea; from -1.2 up to -1.7 °C in Laptev Sea; from -0.8 up to -1.8 °C in northeastern part of Barents Sea. The range of salinity makes from 32 up to 35‰: from 32.3 up to 34.8‰ in Kara Sea, from 26.2 up to

Table 3. Statistical values of basic morphometric characters in Recent *Ciliatocardium ciliatum nordenskiöldi* Kafanov, 1981.

N = 65;  $\bar{y} = L = 18.8 \pm 1.6$ ;  $\bar{y} = 12.8 \pm 1.1$ ;  $As_y = + 0.317$ ;  $E_y = - 0.225$ ;

Parameters	H	A	B	n
x	18.1±1.5	7.8±0.6	5.8±0.5	4.5±0.5
$\sigma_x$	12.4±1.1	5.0±0.4	3.7±0.3	3.7±0.3
$As_x$	+0.315	+0.439	+0.344	+0.236
$E_x$	-0.248	-0.281	-0.210	-0.787
r	0.998±0.000***	0.997±0.001***	0.995±0.001***	0.980±0.004***
$r_{Hx}$	0.999±0.000***	0.998±0.000***	0.996±0.001***	0.980±0.005***
a	1.040	2.122	2.860	6.382
b	0.989±0.005	1.047±0.009	1.080±0.012	0.807±0.020
x/y	0.96±0.00	0.42±0.00	0.32±0.00	0.22±0.01
$\sigma_{xy}$	0.03±0.00	0.02±0.00	0.03±0.00	0.05±0.00
$As_{xy}$	+0.047	+0.147	-0.207	+0.052
$E_{xy}$	-0.380	-0.017	-0.490	-0.387

34.3‰ in Laptev Sea, from 32.4 up to 4.4‰ in northeastern part of Barents Sea. In comparison with Barents Sea in Kara and Laptev Seas it loses its significance in bottom communities.

*Distribution* (Fig. 1): One of the most usual Bivalvia species on the Arctic Ocean shelf: northeastern part of Barents Sea, Kara, Laptev and East Siberian Seas, northwestern part of Chukchi Sea, Beaufort Sea, Canadian Arctic Archipelago, Baffin and Hudson Bays, Labrador Sea.

*Material:* 116 lots (560 specimens): ZIsp.

### *Ciliatocardium ciliatum ochotense* Kafanov,

1981

(Pl. 3, figs. 5-8; pl. 5, figs. 14-18)

*Cardium ciliatum* Fabricius: Ushakov, 1953: 264 (part.);

Scarlato, 1955: 193 (part., non pl. 51, fig. 12).

*Ciliatocardium ciliatum ochotense* Kafanov, 1981a: 54, figs. 3A-G.

*Clinocardium ciliatum* (Fabricius): Volova and Scarlato, 1980: 67, text-fig. 63.

*Bibliography:* Ecology, life cycle, morphology, distribution, fisheries: Kun and Mikulich, 1954; Pasternak, 1957; Saviolov, 1961; Ushakov, 1953; Vinogradov and Neyman, 1969; Vinogradova, 1954; Volova and Scarlato, 1991.

*Etymology:* The name is derived from the type-locality, Okhotsk Sea.

*Type-material:* Holotype (reg. no. 1/200366) and paratypes (reg. nos. 200367-200372) in ZIsp. Type locality: north coast of Okhotsk Sea, west from Tauyskaya Inlet, 58°41' N 147°45' E, depth 121 m, mud; leg. F. A. Derbek, r/v "Okhotsk", September 4, 1912.

Holotype dimensions: L = 48.1; H = 45.0; B = 13.9; A = 18.3.



*Description* (see Table 4): Shell medium-sized, trigonal-suborbicular, high (average H/L ratio about 0.94), sub-equilateral (average A/L ratio about 0.40), medium convex (average B/L ratio about 0.29), faintly oblique and faintly bent, thin-walled, strongly corroded and quite often perforated on the umbonal area. Growth rings smoothed. Umbo low, rather wide, slightly pointed, badly designated by umbonal cavities. Ribs 30-38 (34-35 in average) in number, flattened-triangular in cross section, wrinkled. Longitudinal rows of periostracal fringes on top of ribs are usually present on the anterior and posterior valve areas only. Hinge plate rather narrow. Teeth, especial cardinal, are essentially weakened and frequently are completely reduced. In area of cardinal teeth additional semiinternal ligament is developed.

*Comparison*: The most similar *C. ciliatum nordenskiöldi* Kafanov, 1981 differs by more smoothed shell outlines, by relatively higher, concave-shaped ribs bearing well developed, including on the medial area, longitudinal rows of periostracal fringes.

*Ecology*: It is distributed mainly on mud, sand and sandy mud on depth from 5 up to 174 m but most frequently inhabits depth 50-120 m reflecting preferring to cold intermediate layer of the Okhotsk Sea water mass. In the basic part of area it is found only at constantly negative near-bottom temperatures: from -0.7 up to -1.8 °C. It is usual in *Nuculana pernula* - *Ophiura sarsi* - Polychaeta and *Megayoldia thraciaeformis* - *Macoma calcarea* communities where makes about 6 % (in average about 38.4 g/m<sup>2</sup>) of total zoobenthos biomass.

*Distribution* (Fig. 1): Northern and northwestern portions of Okhotsk Sea.

*Material*: 6 lots (46 specimens): ZIsp.

Table 4. Statistical values of basic morphometric characters in Recent *Ciliatocardium ciliatum ochotense* Kafanov, 1981.

N = 18;  $\bar{y} = L = 37.9 \pm 2.1$ ;  $\bar{y} = 9.0 \pm 1.5$ ;  $As_y = -0.017$ ;  $E_y = +0.755$ ;

Parameters	H	A	B	n
x	35.6±2.1	15.3±0.8	11.1±0.6	9.0±0.7
$\sigma_x$	8.7±1.5	3.5±0.6	2.7±0.5	2.8±0.5
$As_x$	-0.002	-0.056	+0.015	+0.025
$E_x$	+0.424	+0.688	-0.005	+0.093
r	0.996±0.002***	0.968±0.015***	0.986±0.007***	0.969±0.015***
$r_n$	0.997±0.001***	0.980±0.010***	0.989±0.005***	0.986±0.007***
a	1.336	2.659	4.066	8.478
b	0.987±0.020	1.036±0.052	1.014±0.039	0.703±0.030
x/y	0.94±0.00	0.40±0.00	0.29±0.00	0.23±0.01
$\sigma_{x/y}$	0.02±0.00	0.02±0.00	0.01±0.00	0.03±0.01
$As_{x/y}$	-0.025	+0.088	+0.048	-0.032
$E_{x/y}$	+0.139	-0.815	-0.836	+0.310

### *Ciliatocardium ciliatum pubescens* (Couthouy, 1838)

(Pl. 3, figs. 9-12; pl. 4, fig. 5; pl. 6, figs. 7-12)

*Cardium pubescens* Couthouy, 1838: 61, pl. 3, fig. 6.

*Cardium ciliatum* Fabricius: Römer, 1869: 43 (part.), pl. 10, figs. 1, 2 non pl. 7, figs. 5, 6 nec pl. 10, fig. 3; Johnson, 1926: 24 (part.).

*Cardium islandicum* Bruguière: Gould, 1841: 89, fig. 58; DeKay, 1843: 206, pl. 23, fig. 252; Mighels, 1843: 321; Stimpson, 1851: 19; Stimpson, 1854: 21; Gould, 1870: 139, fig. 58; Dall, 1889: 52; Rogers, 1920: 358 (part.).

*Cardium (Cerastoderma) ciliatum* Fabricius: Dall, 1900: 1096 (part.); Dall, 1901: 386 (part.); Whiteaves, 1901: 128 (part.); McLean, 1939: 162 (part.), pl. 24, figs. 1, 2.

*Cardium (Cerastoderma) islandicum* Bruguière: Dautzenberg and Fischer, 1912: 448 (part.).

*Clinocardium ciliatum* (Fabricius): Keen, 1936: 120 (part.); LaRocque, 1953: 63 (part.); Keen, 1954: 20 (part.); Abbott, 1960: 403 (part., an pl. 32, fig. e?); Bousfield, 1960: 28 (part.) non pl. 6, fig. 70.

*Ciliatocardium ciliatum pubescens* (Couthouy, 1838): Kafanov, 1980: 313; 1981: 55, figs. 2G, E.

*Etymology*: The name is derived from the Latin *pubesco*, to be overgrown by hair.

*Type-material*: Most likely it is lost. Dance (1966) marks absence of information on Couthouy's private collection and as probable depositaries names USNM and Museum of Comparative Zoology, Harvard University, Cambridge, where nowadays type-specimens of Couthouy are absent (J. Rosewater, pers. comm., July 17, 1974; K. Boss, pers. comm., 20 October, 1974). Type-locality: [near Egg Rock, off Lynn], Massachusetts Bay, Massachusetts, U.S.A.

Original figure dimensions: L = 20.4; H = 17.8; B = 7.0.

*Description*: Shell medium-sized, orbicular, rather high (average H/L ratio about 0.86) and equilateral (average A/L ratio about 0.44), medium convex (average B/L ratio about 0.34), non-oblique and unbent, thin-walled. Anterior branch of hinge margin inconspicuous, feebly impressed. Posteroventral corner is not expressed. Anterior, ventral and posterior valve margins form a wide correct arch. Growth rings are shown as narrow dark concentric strips. Umbo low, rather wide, very seldom and only slightly corroded. Ribs 36-41 in number, their edges covered with a short, downy ciliation of a dirty brown color. Longitudinal rows of periostracal fringes quite often are absent on the medial valve area however traces of an attachment of these fringes always are kept here. Lunula lengthened and depressed because of what anterior branch of hinge margin is put slightly forward in the commissure plane.

Hinge is a little weakened in comparison with nominative subspecies. Pallial line and muscle scars are poorly distinct because of a strongly brilliant internal surface of valve.

**Comparison:** In shell-shape *C. ciliatum pubescens* strongly differs from all other subspecies and more all reminds *Ciliatocardium likharevi* Kafanov in Scarlato, 1981. However it is not possible to confuse these forms as the last species is absent in North Atlantic.

**Ecology:** On southern and southwestern slopes of Great Newfoundland Bank it is found out on depths 105-247 m at near-bottom temperature in August from + 1.6 up to 2.0 and salinity from 33.98 up to 34.09‰; in Massachusetts Bay on 79m and + 9.5 ; at Cape Cod on depth about 64m.

**Distribution** (Fig. 1): Northeastern Atlantic: from southern and southwestern slopes of Great Newfoundland Bank up to Cape Breton Island, along Nova Scotia coast, Maine Bay, Fundy Bay, south up to Cape Cod, Massachusetts, U.S.A. Dall's (1889) indication for Hatteras Cape, North Carolina, U.S.A. have not proved to be true.

**Material:** 14 lots (36 specimens): Zisp, LACMNH, CAS.

***Ciliatocardium ciliatum tchuktchense* Kafanov in Scarlato, 1981**

(Pl. 4, figs. 1-4; pl. 6, figs. 1-6, 13-15; pl. 7, figs. 1-4)

*Cardium islandicum* Bruguière: Dall, 1871: 148; Krause, 1885: 27; Rogers, 1920: 358 (part.).

*Cardium ciliatum* Fabricius: Leche, 1883: 443 (part.); Scarlato, 1955: 193 (part.) non pl. 51, fig. 12.

*Cardium (Cerastoderma) ciliatum* Fabricius: Dall, 1900: 1096 (part.); Dall, 1901: 390 (part.); Oldroyd, 1925: 142 (part.) non pl. 19, figs. 8, 8a (reproduced from: Sars, 1878, pl. 5, fig. 4; Kuroda and Koba, 1933: 164 (part.); Gorbunov, 1952: 250.

*Cardium (Cerastoderma) islandicum* Bruguière: Dautzenberg and Fischer, 1912: 448 (part.).

*Cerastoderma ciliatum* (Fabricius): Dall, 1921: 39.

*Clinocardium ciliatum* (Fabricius): Keen, 1936: 120 (part.); Habe, 1951: 150 (part.); Morris, 1952: 39, pl. 9, fig. 9; LaRocque, 1953: 63 (part.); Keen, 1954: 20 (part.); Kira, 1954: 111, pl. 55, fig. 2; Kira, 1959: 138, pl. 55, fig. 2; MacGinitie, 1959: 176, pl. 26, fig. 4; Kira, 1962: 156 (part.), pl. 56, fig. 2; Kotaka, 1962: 151 (part.), pl. 35, fig. 11; Habe and Ito, 1965: 132, pl. 44, figs. 3, 4; Petrov, 1966: 221, pl. 17, figs. 1, 2, ?3; Habe and Kosuge, 1967: 148, pl. 55, fig. 21; Abbott, 1968: 228 (part.) non text-fig. 6; Kwon *et al.*, 1993: 364, pl. 85, figs. 3-1, 2.

*Ciliatocardium ciliatum tchuktchense* Kafanov in Scarlato,

1981: 343, photo 310, 311; Kafanov, 1981a: 56, figs. 3D-Z; Volova and Scarlato, 1991: 167 (part.).

*Clinocardium (Ciliatocardium) ciliatum* (Fabricius): Higo and Goto, 1993: 617 (part.); Coan and Scott, 1997: 13; Higo *et al.*, 1999: 474 (part.).

**Bibliography:** Ecology, life cycle, morphology, distribution, fisheries: Bernard, 1967, 1979; Buyanovsky, 1994; Dall, 1919; Evseev, 1981; Eyerdam, 1960; Feder and Jewett, 1981; Feder *et al.*, 1994; Filatova and Barsanova, 1964; Galkin, 1953; Gordeeva, 1948; Habe and Igarashi, 1967; Ito, 1967; Kafanov, 1978a, b, 1979, 1981b; Keen, 1937; Klimova, 1974; Kulichkova, 1955; Kun and Mikulich, 1954; Kuroda and Habe, 1952; Kuroda and Kinoshita, 1951; Kuznetsov, 1961, 1963, 1970; MacGinitie, 1955; McLaughlin, 1963; Neyman, 1960; Pasternak, 1957; Saviolov, 1961; Scalkin, 1963; Vinogradov and Neyman, 1969; Vinogradova, 1954; Zenkevich, 1963.

**Etymology:** The name is derived from the part of area, Chukchi Sea.

**Type-material:** Holotype (reg. no. 1/200291) and paratypes (reg. nos. 200292-200365) in ZIsp. Type locality: Bering Sea, Provideniya Bight, 67° 25' N 173° 10' W, depth 22-11 m, mud; leg. E. E. Arngold, r/v "Vaygach", August 6 (19), 1912.

Holotype dimensions: L = 54.9; H = 52.9; B = 16.4; A = 23.3.

**Description** (see Table 5): Shell medium-sized, angular-subcircular or angular-suborbicular, high (average H/L ratio about 0.95), slightly subequilateral (average A/L ratio about 0.43), medium convex (average B/L ratio about 0.30), faintly oblique and faintly bent, rather thick-walled. Almost straight posterior branch of hinge margin passes in posterior margin of valve with an appreciable corner; posteroventral corner is well expressed and usually situated down- or backward. Growth rings well marked. Umbo

Table 5. Statistical values of basic morphometric characters in Recent *Ciliatocardium ciliatum tchuktchense* Kafanov in Scarlato, 1981.

N = 71;  $\bar{y} = L = 34.7 \pm 1.4$ ;  $\bar{y}_y = 12.1 \pm 1.0$ ;  $As_y = +0.146$ ;  $E_y = +0.140$ ;

Parameters	H	A	B
x	33.0±1.4	14.6±0.5	10.3±0.4
$\sigma_x$	11.7±1.0	4.6±0.4	3.7±0.3
$As_x$	+0.229	+0.117	+0.291
$E_x$	+0.522	+0.118	+0.343
r	0.995±0.001***	0.992±0.002***	0.985±0.004***
$r_{Hx}$	0.997±0.001***	0.993±0.002***	0.990±0.002***
a	1.128	1.702	3.828
b	0.988±0.009	1.093±0.015	1.010±0.017
x/y	0.95±0.00	0.43±0.00	0.30±0.00
$\sigma_{x/y}$	0.03±0.00	0.03±0.00	0.02±0.00
$As_{x/y}$	-0.358	+0.272	+0.180
$E_{x/y}$	+0.091	+0.522	-0.661

marked mainly by anterior umbonal cavity, wide and medium height, often slightly corroded. Ribs 30-38 (33-35 in average) in number, rather low and wide. Ciliated periostracal fringes on the ribs are poorly advanced and mainly on the posterior area only. As a rule, the hinge is not reduced. Lunula rather small, lengthened, slightly pressed.

*Comparison:* From the most similar *C. ciliatum ciliatum* it differs by more angular outlines of shell and by relatively more equilateral valves.

*Ecology:* Off eastern Kamchatka and northern Kurile Islands it meets on depths 40-445 m where often forms *C. ciliatum* community and gives a density 1-16 (3 in average) specimens/m<sup>2</sup> and biomass 0.3-115.0 (41.3 in average) g/m<sup>2</sup> at close to 0 near-bottom temperature and salinity 33.0-33.5‰.

*Distribution* (Fig. 1): From East Korea Bay, Japan Sea, Tajima, Hyogo Prefecture, Japan, Otsuchi Bay, northeastern Honshu, Japan, Commander and Aleutian (west to Amchitka Island) Islands, east Kodiak Island, Alaska, Puget Sound, Washington, U.S.A., north- and eastward up to southeastern part of Chukchi Sea and Point Barrow, Arctic coast of Alaska.

*Material:* 96 lots (more than 550 specimens): ZIsp, LACMNH, CAS, SBMNH.

### ***Ciliatocardium ciliatum chikagawaense***

(Kotaka, 1950)

(Pl. 7, figs. 5, 6)

*Cardium shinjiense* "Yokoyama, 1923": Yokoyama, 1926, pl. 34, figs. 17, 18.

*Clinocardium chikagawaense* Kotaka, 1950: 46, pl. 5, figs. 1-6; Ogasawara, 1977: 114, pl. 11, fig. 15; Iwai, 1960, pl. 1, fig. 10; Hatai *et al.*, 1961, pl. 2, fig. 9; Kanno, 1962, pl. 5, fig. 6; Iwai, 1965: 38, pl. 17, figs. 18a-b, 20.

*Ciliatocardium ciliatum chikagawaense* (Kotaka): Kafanov, 1980: 312.

? *Clinocardium chikagawaense* Kotaka, 1950: Matsui, 1985: 231, pl. 33, fig. 11.

*Clinocardium ciliatum* (Fabr.): Ogasawara *et al.*, 1986, pl. 41, figs. 73, 4; pl. 62, figs. 11a,b.

Non

*Clinocardium chikagawaense* Kotaka, 1950: Shimamoto and Koike, 1986: 38, pl. 5, figs. 1, 4, 6, 8a-b; pl. 6, fig. 10.

*Etymology:* The name is derived from the type-locality, Chikagawa.

*Type-material:* Holotype - IGPS no. 72999. Type-locality: the sea cliff at the outlet of Chikagawa River at Chikagawa, Tanabu-machi, Shimokita District, Aomori

Prefecture, Honshu, Japan; Pliocene Hamada Formation.

Holotype dimensions: L = 58.4; H = 58.0; B = 21.0; A = 24.4.

*Description:* Shell medium-sized, trigonal-suborbicular, high or extended in height (average H/L ratio about 0.99), subequilateral (average A/L ratio about 0.42), convex (average B/L ratio about 0.36), faintly oblique and faintly bent, rather thick-walled. Postero-dorsal margin broadly arched, long, extending to prolonged posterior margin, than sharply merging into rather narrowly rounded ventral margin. Antero-dorsal margin not shouldered, roundly protruding into anterior margin. Angle of antero- and postero-dorsal margins at umbo is about 114°. Growth rings are distinct in later stage of growth but undistinct in young stages. Umbo tapering, moderate high, markedly incurved and directed anteriorly. Ribs 30-40 in number, wrinkled, crossed by numerous fine concentric growth lines. Ribs mostly straight, sharp edged, broader than flatt-bottomed intercostal spaces sometimes shallowly excavated. Ligament area low, and narrow. Lunule circumscribed and heart-shaped. Cardinal teeth situated nearer to anterior laterals than to posterior. Recurved tooth *3b* stronger than *3a*. The tooth *AI* much stronger than *PI*. Muscle scars distinct, anterior one more oval than quadratic posterior one.

*Comparison:* This subspecies resembles Recent *Ciliatocardium ciliatum* (Fabricius, 1780) in many respects but can be distinguished by the more angularly or postero-ventrally elongated shell outline and by markedly incurved and directed anteriorly umbo.

*Distribution:* It is one of the most characteristic representatives of the Pliocene so-called Omma fauna in North Japan. Honshu: Hamada, Narusawa and Daishaka Formations of Aomori Prefecture; Sawane Formation of Niigata Prefecture; Shibikawa and Sasaoka Formation of the Akita oil-field. Setana Formation of Hokkaido.

### ***Ciliatocardium ciliatum maruyamense* Kafanov et Savizky, 1982**

(Pl. 7, figs. 7, 8)

? *Laevicardium (Cerastoderma) shinjiense* (Yok.): Ilyina, 1954 [1957]: 223, pl. 16, figs. 6, 7, 8, 8a.

*Ciliatocardium ciliatum maruyamense* Kafanov and Savizky, 1982: 56, pl. 5, figs. 4, 5.

Non

*Ciliatocardium ciliatum marujamensis* Kafanov: Gladenkov *et al.*, 1992, pl. 3, fig. 8.

*Etymology:* The name is derived from the type-locality, Maruyamskaya Suite.

*Type-material:* Holotype - PIN no. 28/3962. Type-locality: Malyi Takoy River, Dolinsk District, Sakhalin, Russia; lower sub-suite of Maruyamskaya Suite (stratotypical section), upper Miocene.

Holotype dimensions: L = 43.2; H = 41.0; B = 12.5; A = 15.7.

*Description* (see Table 6): Shell medium-sized, trigonal-suborbicular, high (average H/L ratio about 0.96), sub-equilateral (average A/L ratio about 0.42), medium convex (average B/L ratio about 0.31), faintly oblique and faintly bent, rather thick-walled. Growth rings smoothed. Umbo medium height and medium wide, slightly marked by anterior umbonal cavity only. Ribs 28-35 in number (more often 29-32), rather wide and low, appreciably bent on all external surface of valve. On anterior area ribs are a little placed, on posterior one pulled together and smoothed. Intercoastal spaces are some narrower than ribs.

*Comparison:* From Recent North Pacific *C. ciliatum tchuktchense* Kafanov in Scarlato, 1981 it differs by relatively lesser average ribs number (29-32 against 33-35) and by main conchometric parameters in regressions of H, B and A against L (compare Tables 5 and 6). Paired comparison of regression coefficients shows significant statistical differences at highest significance level ( $P = 0.999$ ). *Ciliatocardium ciliatum chikagawaense* well distinguishes by much higher shell.

*Distribution:* Upper Miocene lower-subsuite of Maruyamskaya Suite in South Sakhalin.

*Material:* 16 lots (32 specimens): GIN, VNIGRI, MGRI.

Table 6. Statistical values of basic morphometric characters in *Ciliatocardium ciliatum maruyamense* Kafanov et Savizky, 1982.

$N = 25$ ;  $y = L = 40.7 \pm 0.9$ ;  $y = 4.6 \pm 0.6$ ;  $As_y = -0.536$ ;  $E_y = -0.108$ ;

Parameters	H	A	B	B
$\bar{x}$	39.2±1.1	17.0±0.4	12.5±0.3	11.7±0.4
$\sigma_x$	5.4±0.8	1.8±0.3	1.6±0.2	2.1±0.3
$As_x$	-0.462	-0.132	-0.438	+0.399
$E_x$	-0.648	-0.084	+0.499	-0.717
$r$	0.994±0.001***	0.990±0.002***	0.985±0.004***	0.983±0.005***
$r_{12}$	0.997±0.001***	0.993±0.002***	0.990±0.002***	0.990±0.003***
$a$	2.460	2.776	5.551	10.402
$b$	0.765±0.060	0.948±0.117	0.789±0.079	0.554±0.082
$x/y$	0.96±0.01	0.42±0.00	0.31±0.00	0.29±0.01
$\sigma_{x/y}$	0.04±0.01	0.02±0.00	0.01±0.00	0.03±0.00
$As_{x/y}$	+0.153	+0.343	-0.187	+0.352
$E_{x/y}$	-0.394	-0.749	+0.645	-0.203

### *Ciliatocardium likharevi* Kafanov in Scarlato,

1981

(Pl. 7, figs. 9-16)

*Cardium ciliatum* (Fabricius): Scarlato, 1955, pl. 51, fig. 12.

*Ciliatocardium likharevi* Kafanov: Scarlato, 1981: 344, photo 312; Kafanov, 1981a: 58, text-figs. 1A; 4A-G.

*Clinocardium (Ciliatocardium) ciliatum* (Fabricius): Higo and Goto, 1993: 617 (part.).

Non

*Ciliatocardium likharevi* Kafanov.: Volova and Scarlato, 1991: 167.

*Etymology:* The name is derived in the honour of well-known Russian malacologist Prof. Iliya M. Likharev (Zoological Institute, Russian Academy of Sciences).

*Type-material:* Holotype- ZIsp no. 1/200274. Type-locality: Japan Sea, Tatar Strait, off Rakuma (now Antonovo Village), Sakhalin, Russia, depth 45-40 m, sand.

Holotype dimensions: L = 43.2; H = 41.0; B = 12.5; A = 15.7.

*Description* (see Table 7): Shell medium-sized, suborbicular or ellipsoid, high (average H/L ratio about 0.90), equilateral (average A/L ratio about 0.44), medium convex (average B/L ratio about 0.30), non-oblique and unbent, rather thick-walled. Short anterior branch of hinge margin is clearly concave and is put forward to the valves commissure plane. Growth rings usually smoothed. Umbo tapering, moderate wide, well marked at least by anterior umbonal cavity. Ribs 33-40 (36-37 in average) in number, rather low, with almost straight shape of slope. Three most forward ribs are shortened, strong flattened and in cross section have almost rounded form. Ribs covered by numerous concentric periostracal wrinkles. Hinge plate rather wide. The teeth *AI* and *AIII* misses under some corner, the last one is considerably weakened. The teeth 2 and 4b settle down to each other under a corner approximately 90°. The straight line nymph is equal to length of a tooth *PI*. Lunula precise, lanceolate.

*Comparison:* From *C. ciliatum* (Fabricius, 1780) it differs by equilateral, non-oblique and unbent shell, by concave anterior branch of dorsal margin and by relatively shorter nymph (in average no more than 1/5 of the shell length).

*Taxonomical notes:* The taxonomic independence of this true sibling species is strictly proved by the statistical analysis at its joint sites with *C. ciliatum tchuktchense* (see Kafanov, 1981a).

*Ecology:* It meets on depths from 33 m (Aniwa Bay) up

Table 7. Statistical values of basic morphometric characters in *Ciliatocardium likharevi* Kafanov in Scarlato, 1981.

N = 25; y = L = 20.7 ± 2.1;  $\bar{y}$  = 9.8 ± 1.5;  $As_y$  = + 0.411;  $E_y$  = + 0.629;

Parameters	H	A	B	n
x	18.7±2.0	9.1±0.9	6.1±0.6	3.8±0.5
$\sigma_x$	8.9±1.4	4.2±0.6	2.8±0.4	2.1±0.3
$As_x$	+0.357	+0.369	+0.364	+0.481
$E_x$	+0.447	+0.495	+0.575	+0.562
r	0.999±0.000***	0.995±0.002***	0.994±0.003***	0.981±0.008***
$r_{ln}$	0.999±0.000***	0.993±0.003***	0.988±0.005***	0.969±0.013***
n	1.080	2.244	3.388	6.664
b	0.976±0.012	1.018±0.029	1.023±0.037	0.818±0.048
x/y	0.90±0.00	0.44±0.00	0.30±0.00	0.18±0.00
$\sigma_{x/y}$	0.02±0.00	0.02±0.00	0.02±0.00	0.02±0.00
$As_{x/y}$	+0.378	-0.208	+0.201	-0.112
$E_{x/y}$	-0.472	-0.055	-0.377	-0.295

to 117 m (South Kurile shallow-water) on mud and, less often, on sand at August near-bottom temperature from -0.3 (Aniwa Bay) up to 12.8 (South Kurile shallow-water).

**Distribution:** Japan Sea: off south Primorye Territory, including Peter the Great Bay, southwestern Sakhalin, Russia; Okhotsk Sea: Aniwa and Terpeniya Bays; South Kurile shallow-water. Fresh empty valves are found off Nagasaki, Kyushu.

**Material:** 22 lots (26 specimens): ZIsp, Museum of Institute of Marine Biology (Vladivostok City).

## References

- Abbott, R. T. (1960), *American sea-shells*. 5<sup>th</sup> printing. 541 pp., 40 pls. D. Van Nostrand, Princeton, New Jersey, etc.
- Abbott, R. T. (1968), *Seashells of North America; A guide to field identification*. 280 pp. Golden Press, New York.
- Adams, H. and Adams, A. (1858), *The genera of Recent Mollusca; arranged according to their organization*. Vol. 2. 660 pp. John van Voorst, London.
- Allen, J. A. (1965), Records of Mollusca from the northwest Atlantic obtained by Canadian fishery research vessels, 1946-61. *J. Fish. Res. Bd. Canada*, **22**(4), 977-997.
- Andrews, J. T. (1972), Recent and fossil growth rates of marine bivalves, Canadian Arctic, and Late-Quaternary Arctic marine environments. *Palaeogeogr., Palaeoclimatol., Palaeoecol.*, **11**(3), 157-176.
- Bernard, F. R. (1967), Prodrôme for a distributional check-list and bibliography of the Recent marine Mollusca of the west coast of Canada. *Techn. Rept. Fish. Res. Bd. Canada*, no. 2, i-xxiv, 1-261.
- Bernard, F. R. (1979), Bivalve mollusks of the western Beaufort Sea. *Nat. Hist. Mus. Los Angeles County, Contribs. Sci.*, no. 313, 1-80.
- Bousfield, E. L. (1960), *Canadian Atlantic sea shells*. 72 pp., 13 pls. Natl. Mus. Canada, Ottawa.
- Broderip, W. J. and Sowerby, G. B. I. (1829), Observations on new or interesting Mollusca contained, for the most part, in the Museum of the Zoological Society. *Zool. J.*, **4**(15), 359-376, pl. 9.
- Brøgger, W. C. (1901), Om de sennglaciale og postglaciale nivåforandringer i Kristianiafeltet. *Norges Geol. Unders.*, no. 31, 67-82.
- Brotskaya, V. A. and Zenkevich, L. A. (1939), The quantitative account of the Barents Sea bottom fauna. *Transact. All-Union Res. Inst. Fish. Oceanogr. (VNIRO)*, **4**, 5-126 (in Russian).
- Brotskaya, V. A., Zhdanova, N. N. and Semenova, N. L. (1963), Bottom fauna of Velikaya Salma and adjacent areas of Kandalaksha Bay of the White Sea. *Transact. Kandalaksha State Reserve*, **4**, 159-182 (in Russian).
- Bruguère, J. G. (1789), *Encyclopédie Méthodique; histoire naturelle des vers, des mollusques, des coquillages et zoophytes*. In: *Histoire Naturelle des Vers*. **1**(1). 344 pp. Panckoucke, Paris.
- Buyanovskiy, A. I. (1994), *Marine bivalve molluscs of Kamchatka and perspectives of their management*. 100 pp. All-Russ. Res. Inst. Fish. Oceanogr. (VNIRO), Moscow (in Russian).
- Chemnitz, J. H. (1782), *Neues systematisches Conchylien-Cabinet*. Vol. 6. 375 S., 36 Tafn. G. N. Raspe, Nürnberg.
- Chinzei, K. (1959), Molluscan fauna of the Pliocene Sannohe group of Northeast Honshu, Japan. 1. The faunule of the Kubo formation. *J. Fac. Sci., Univ. Tokyo, Ser. 2*, **12**(1), 103-132, pls. 9-11.
- Clarke, A. H. (1974), Molluscs from Baffin Bay and the northern North Atlantic Ocean. *Natl. Mus. Canada Publ. Biol. Oceanogr.*, no. 7, 1-23.
- Clarke, A. H., Grant, D. R. and Macpherson, E. (1972), The relationship of *Atractodon stonei* (Pilsbry) (Mollusca, Buccinidae) to the Pleistocene stratigraphy and paleo-ecology of Southwestern Nova Scotia. *Canad. J. Earth Sci.*, **9**(8), 1030-1038.
- Clench, W. J. and Smith, L. C. (1944), The family Cardiidae in the Western Atlantic. *Johnsonia*, **1**(13), 1-32, pls. 1-13.
- Coan, E. V. and Scott, P. H. (1997), Checklist of the marine bivalves of the Northeastern Pacific Ocean. *Santa Barbara Mus. Nat. Hist., Contribs. Sci.*, **1**, 1-28.
- Couthouy, J. P. (1838), Descriptions of new species of Mollusca and shells, and remarks on several polypi, found in Massachusetts Bay. *Boston J. Nat. Hist.*, **2**(1), 53-111, pls. 1-3.
- Dall, W. H. (1871), Descriptions of sixty new forms of mollusks from the west coast of North America and the North Pacific Ocean, with notes on other already described. *Amer. J. Conchol.*, **7**(2), 93-160, pls. 13-16.
- Dall, W. H. (1889), A preliminary catalogue of the shell-bearing marine mollusks and brachiopods of the southeastern coast of the United States, with illustrations of many of the species. *Bull. U. S. Natl. Mus.*, no. 37, 1-221, pls. 1-74.
- Dall, W. H. (1900), Contributions to the Tertiary fauna of Florida, with especial reference to the Miocene silex-beds of Tampa and the Pliocene beds of the Caloosahatchie River, including in many cases a complete revision of the generic

- groups treated of and their American Tertiary species. Pt. V. Teleodermacea: *Solen to Diplodonta*. *Transact. Wagner Free Inst. Sci.*, **3** (5), 949-1218, pls. 36-47.
- Dall, W. H. (1901), Synopsis of the family Cardiidae and of the North American species. *Proc. U. S. Natl. Mus.*, **23** (1214), 381-392.
- Dall, W. H. (1919), The Mollusca of the Arctic coast of America collected by the Canadian Arctic expedition west from Bathurst Inlet with an appended report on a collection of Pleistocene fossil Mollusca. *Repts. Canad. Arctic Exped., 1913-1918*, **8A**, 3-23, 26-29, pls. 1-3.
- Dall, W. H. (1921), Summary of the marine shell-bearing mollusks of the northwest coast of America, from San Diego, California, to the Polar Sea, mostly contained in the collection of the United States National Museum, with illustrations of hitherto unfigured species. *Bull. U. S. Natl. Mus.*, no. 112, iii + 217 pp., 22 pls.
- Dall, W. H. (1924a), Report on the Pleistocene fossils collected on the Arctic coast. *Repts. Canad. Arctic Exped., 1913-1918*, **11A**, 30-33.
- Dall, W. H. (1924b), List of mollusks from the eastern Arctic region, collected by the Neptune expeditions, etc. *Repts. Canad. Arctic Exped., 1913-1918*, **8A** (suppl. Report), 33-35.
- Dance, S. P. (1966), *Shell collecting, an illustrated history*. 344 pp. Univ. California Press, Berkeley & Los Angeles.
- Dautzenberg, Ph. and Fischer, H. (1912), Mollusques provenant des campagnes de l'Hirondelle et de la Princesse-Alice dans les mers du nord. *Rés. Camp. Sci. Accompl. Yacht Albert 1er Prince Souv. Monaco*, Fasc. 37, 1-629, pls. 1-11.
- DeKay, J. E. (1843), *Natural history of New York. Pt. 5. Mollusca*. 277 pp., 40 pls. Published by the State, Albany.
- Derjugin, K. M. (1915), Fauna of the Kolsky Bay and condition of its existence. *Méms. L'Acad. Impér. Sci. St.-Petersbourg. Sci. Mathém., Physiq. et Natur.* Ser. 8, **34** (1), 1-929 (in Russian).
- Derjugin, K. M. (1928), Fauna of the White Sea and condition of its existence. *State Hydrol. Inst., Seas Res.*, Issue 7-8, 1-511 (in Russian).
- Durham, J. W. and MacNeil, F. S. (1967), Cenozoic migrations of marine invertebrates through the Bering Strait region. In Hopkins, D. M. (ed.), *The Bering land bridge*. 326-349. Stanford University Press.
- Evseev, G. A. (1981), *Communities of bivalve molluscs in the postglacial deposits of the Sea of Japan shelf*. 160 pp., 8 pls. Nauka, Moscow. (in Russian).
- Eyerdam, W. J. (1960), Mollusks and brachiopods from Afognak and Sitkalidak Islands, Kodiak Group, Alaska. *The Nautilus*, **74** (2), 41-46, 46a-46e.
- Fabricius, O. (1780), *Fauna Groenlandica, systematice sistens Animalia Groenlandiae occidentalis hactenus indagata, etc.* xvi + 452 pp., 1 pl. Rothe, Hafniae & Lipsiae.
- Feder, H. M., Foster, N. R., Jewett, S. C., Weingartner, T. J. and Baxter, R. (1994), Mollusks in the Northeastern Chukchi Sea. *Arctic*, **47** (2), 145-163.
- Feder, H. M. and Jewett, S. C. (1981), Feeding interactions in the Eastern Bering Sea with emphasis on the benthos. In Hood, D. W. and Calder, J. A. (eds.), *The Eastern Bering Sea shelf: oceanography and resources. Vol. 2*. 1229-1261. U. S. Dept. Commerce, Washington, D. C.
- Fedyakov, V. V. (1986), *Molluscs of the White Sea... Objective conformities of distribution*. 127 pp. Zool. Inst., Acad. Sci. USSR, Leningrad (in Russian).
- Filatova, Z. A. (1948), Class bivalve molluscs (Bivalvia, Lamellibranchiata). In Gaevskaya, N. S. (ed.), *A guide to fauna and flora of the northern seas of the USSR*. 405-446, pls. 105-113. Sovetskaya nauka, Moscow (in Russian).
- Filatova, Z. A. and Barsanova, N. G. (1964), Communities of bottom fauna in western part of Bering Sea. *Transact. Inst. Oceanol., Acad. Sci. USSR*, **69**, 6-97 (in Russian).
- Filatova, Z. A. (1957), An outlook on bivalve molluscs fauna of the northern seas of the USSR. *Transact. Inst. Oceanol., Acad. Sci. USSR*, **20**, 3-59 (in Russian).
- Fischer-Piette, E. (1977), Revision des Cardiidae (mollusques lamellibranches). *Méms. Mus. natl. d'Hist. natur., Nouv. Ser.*, A, Zoologie, **101**, 1-212, pls. 1-12.
- Friele, H. and Grieg, J. A. (1901), *Mollusca. III. The Norwegian North-Atlantic Expedition, 1876-1878, Zoology*. viii+130 pp. Grøndahl a. Søn, Christiania.
- Galkin, Yu. (1953), Materials to the characteristic of molluscan fauna of Aniwa Bay (Sea of Okhotsk). *Transact. Zool. Inst., Acad. Sci. USSR*, **13**, 242-249 (in Russian).
- Gladenkov, Yu. B. (1972), Neogene of Kamchatka (problems of biostratigraphy and paleoecology). *Transact. Geol. Inst., Acad. Sci. USSR*, **214**, 1-252, pls. 1-8 (in Russian with English title).
- Gladenkov, Yu. B., Barinov, K. B., Basilyan A. E., Bordunov, S. I., Bratzeva, G. M., Zyryanov, E. B., Kuralenko, N. P., Vitukhin D. I., Oreshkina, T. V., Ganzey, S. S., Kiyashko, S. I. and Trubikhin, V. M. (1992), Detailed division of the Neogene of Kamchatka. *Transact. Geol. Inst., Russ. Acad. Sci.*, **478**, 1-208 pp., pls. 1-60 (in Russian).
- Gladenkov, Yu. B., Norton, P. and Spaink, G. (1980), Upper Cenozoic of Iceland (Stratigraphy of Pliocene-Pleistocene and paleontological assemblages). *Transact. Geol. Inst., Acad. Sci. USSR*, **345**, 1-116, pls. 1-16 (in Russian).
- Gmelin, J. F. (1791), *Caroli a Linné ... Systema naturae per regna tria naturae ... editio decima tertia, acuta, reformata*. **1** (6), 3021-3910. Beer, Leipzig.
- Golikov, A. N. and Scarlato, O. A. (1977), Composition, distribution and ecology of gastropods and bivalves off Franz Josef Land. In Golikov, A. N. (ed.), *Biocoenoses of the shelf of Franz Josef Land and the Fauna of adjacent waters*. 313-390 Nauka, Leningrad (in Russian).
- Gorbunov, G. P. (1946), Bottom population of Novosibirsk shallow-waters and central part of the Northern Ice Ocean. In Gorbunov, G. P. and Ushakov, P. V. (eds.), *Transactions of the Drifting expedition of Glavsevmorput' on the icebreaking steamship "G. Sedov", 1937-1940. Vol. 3. Biology*. 30-138, pl. 1. Glavsevmorput' Publ. House, Moscow & Leningrad (in Russian).
- Gorbunov, G. P. (1952), Bivalve molluscs (Bivalvia) of the Chukchi Sea and Bering Strait. In Ushakov, P. V. (ed.), *The*

- Extreme North-East of the USSR. Vol. 2. Fauna and flora of the Chukchi Sea.* 216-278. Acad. Sci. USSR Publ. House (in Russian).
- Gordeeva, K. T. (1948), Materials on quantitative study of zoobenthos in the west Kamchatka shelf. *Proc. Pac. Res. Inst. Fish. Oceanogr. (TINRO)*, **26**, 131-198 (in Russian).
- Gould, A. A. (1841), *Report on the Invertebrata of Massachusetts, comprising the Mollusca, Crustacea, Annelida, and Radiata.* 373 pp., 15 pls. (unnumbered, comprising 213 figs.). Cambridge Univ. Press.
- Gould, A. A. (1870), *Report on the Invertebrata of Massachusetts, published agreeably to an order of the legislature. Second edition, comprising the Mollusca.* Ed. by W. G. Binney. 524 pp., pls. 16-27. Wright & Potter, Boston.
- Grant, U. S. IV and Gale, H. R. (1931), Catalogue of the marine Pliocene and Pleistocene Mollusca of California. *Mems. San Diego Soc. Nat. Hist.* **1**, 1-1036, pls. 1-32.
- Grieg, J. A. (1924), Molluscs, brachiopods and echinoderms from Novaya Zemlya. *Rept. Sci. Results Norweg. Exped. Nov. Zemlya 1921*, no. 26, 1-16.
- Grigorenko, Yu. N., Krishtofovich, L.V., Tarasov, B. M., Motylinskaya, T. A., Shakhmundes, V. A. and Budasheva, A. I. (1967), Lithological-and-facial, geochemical and paleontological characteristics of stratotypical section of Kavranskaya Series in West Kamchatka. *Transact. VNIGRI*, **254**, 27-114 (in Russian).
- Habe, T. (1951), *Genera of Japanese shells. Pelecypoda, no. 2.* 95-186. Kairui Bunken Kankokai, Kyoto.
- Habe, T. and Igarashi, T. (1967), A list of marine molluscan shells in the Fisheries Museum, Faculty of Fisheries, Hokkaido University (Kawasaki collection and specimens collected by the Marine Zoological Laboratory). *Contribs. Fish. Mus., Fac. Fish., Hokkaido Univ.*, no. 6, 1-56 (in Japanese).
- Habe, T. and Ito, K. (1965), *Shells of the world in colour. Vol. 1. The Northern Pacific.* vii + 176 pp., 56 pls. Hoikusha, Osaka.
- Habe, T. and Kosuge, S. (1967), *The standard book of Japanese shells illustrated in color.* xviii + 223 pp., 64 pls. Hoikusha, Osaka (in Japanese).
- Hägg, R. (1905), Mollusca und Brachiopoda gesammelt von der schwedischen zoologischen Polarexpedition nach Spitzbergen, dem nordöstlichen Grönland und Jan Mayen im J. 1900, 1. *Ark. Zool.*, **2**(2), 1-66, Taf. 1.
- Hancock, A. (1846), A list of shells dredged on the west coast of Davis's Strait; with notes and descriptions of eight new species. *Anns. Mag. Nat. Hist.*, **18** (120), 323-338, pl. 5.
- Hatai, K., Masuda, K. and Suzuki, Y. (1961), A note on the Pliocene megafossil fauna from the Shimokita Peninsula, Aomori Prefecture, Northeast Honshu, Japan. *Saito Ho-on Kai Mus., Res. Bull.*, no. 30, 18-38, pls. 1-4.
- Herzenstein, S. (1885), Materials to the fauna of Murman Coast and White Sea. I. Molluscs. *Transact. St.-Petersburg Soc. Naturalists*, **16**, 635-814 (in Russian).
- Higo, S., Callomon, P. and Goto, Y. (1999), *Catalogue and bibliography of the marine shell-bearing Mollusca of Japan. Gastropoda, Bivalvia, Polyplacophora, Scaphopoda.* 749 pp. Elle Sci. Publ., Osaka.
- Higo, S. and Goto, Y. (1993), *A systematic list of molluscan shells from the Japanese Is. and the adjacent area.* 693+148 pp. Yuko, Osaka (in Japanese).
- Hopkins, D. M. (1959), Cenozoic history of the Bering land bridge. *Science*, **129**(3362), 1519-1528.
- Hopkins, D. M., MacNeil, F. S., Merklin, R. L. and Petrov, O. M. (1965), Quaternary correlations across Bering Strait. *Science*, **147**(3662), 1107-1114.
- Ilyina, A. P. (1954 [published under the griff "top secret"; data of the opening for public use is May 12, 1957]), Molluscs of the Neogene deposits of South Sakhalin. *Transact. VNIGRI, Spec. Ser.*, fasc. 10, 188-314, pls. 1-30 (in Russian).
- Ilyina, A. P. (1960), Molluscs of the Tertiary deposits of Koryak mountain ridge. *Transact. VNIGRI, fasc. 154, Paleontological collection*, 2, 306-313, pls. 1-2 (in Russian).
- Ito, K. (1967), A catalogue of the marine molluscan shell-fish collected on the coast of and off Tajima, Hyogo Prefecture. *Bull. Japan Sea Reg. Fish. Res. Lab.*, no. 18, 39-91, pls. 1-10.
- Iwai, T. (1960), Pliocene Mollusca from the Nishi-Tsugaru district, Aomori Prefecture, Japan. *Saito Ho-on Kai Mus., Res. Bull.*, no. 29, 35-46, pls. 1-4.
- Iwai, T. (1965), The geological and paleontological studies in the marginal area of the Tsugaru basin, Aomori Prefecture, Japan. *Bull. Educ. Fac., Hirosaki Univ.*, no. 15, 1-68, pls. 12-20.
- Iwai, T. and Siobara, T. (1969), Pleistocene Mollusca from Kamikita-gun, Aomori Prefecture, Japan. *Bull. Educ. Fac., Hirosaki Univ.*, no. 20, 1-7, pls. 1-3.
- Jeffreys, J. G. (1864), *British conchology, or an account of the Mollusca which now inhabit the British Isles and the surrounding seas. Vol. II. Marine shells, comprising the Brachiopoda, and Conchifera from the family Anomiidae to that of the Macrtridae.* xiv + 566 pp., 8 pls. John van Voorst, London.
- Jensen, A. S. (1901), Om levninger af grundtvandsdyr på store havdyb mellem Jan Mayen og Island. *Vidensk. Medd. Dansk Naturhist. Forening*, **52**, 229-239.
- Jensen, A. S. (1912), *Lamellibranchiata. Part I. The Danish Ingolf-Expedition. Vol. 2, Pt. 5.* 119 pp., pls. 1-4. H. Hagerup, Copenhagen.
- Johnson, C. W. (1926), A list of the molluscs collected by Mr. Owen Bryant along the coast of Labrador, New Foundland and Nova Scotia. *Nautilus*, **39**, 128-135; **40**, 21-25.
- Kafanov, A. I. (1974), Composition, taxonomy and evolution of the group *Clinocardium* (Mollusca, Cardiidae). *Zool. Zh. [Zool. J.]*, **53**(10), 1466-1476 (in Russian with English summary).
- Kafanov, A. I. (1978a), Temperature variability of linear growth and life duration in six species of the subfamily Clinocardiinae Kafanov, 1975 (Mollusca, Cardiidae). *Zool. Zh. [Zool. J.]*, **57**(10), 1480-1488 (in Russian with English summary).
- Kafanov, A. I. (1978b), Some aspects of ecological interrelationship between three species of the subfamily Clinocardiinae (Bivalvia, Cardiidae). In Golikov, A. N. (ed.), *Regularities of distribution and Ecology: of coastal marine biocenosis:*

- Materials of the USSR-USA Symposium on the Program "Biological productivity and biochemistry of the World's Ocean. 86-88 (in Russian), 216-218 (in English). Nauka, Leningrad.*
- Kafanov, A. I. (1979), On conservatism and variability in growth temperatures of marine molluscan shells. *Biologiya Morya (Marine Biology, Vladivostok)*, no. 6, 59-69 (in Russian with English summary).
- Kafanov, A. I. (1980), Systematics of the subfamily Clinocardiinae Kafanov, 1975 (Bivalvia, Cardiidae). *Malacologia*, **19**(2), 297-328.
- Kafanov, A. I. (1981a). A revision of the genus *Ciliatocardium* Kafanov, 1974 (Bivalvia, Cardiidae). In Kafanov, A. I. (ed.), *The systematics and chorology of bottom invertebrates from the Far East seas*. 43-61, pls. 2-3. Far East Sci. Center, Acad. Sci. USSR, Vladivostok (in Russian).
- Kafanov, A. I. (1981b). Determination of paleotemperatures in Neogene sea basins by duration of marine molluscs' life. *Biologiya Morya (Marine Biology, Vladivostok)*, no. 5, 15-20 (in Russian with English summary).
- Kafanov, A. I. and Savizky, V. O. (1982), A review of Paleogene and Neogene *Ciliatocardium* (Bivalvia Cardiidae) of Sakhalin. *Paleontol. Zh. (Paleontol. J.)*, no. 3, 53-61. (in Russian).
- Kanehara, K. (1937), On *Cardium (Cerastoderma) shinjiense* Yokoyama, with description of a new species, *C. uyemurai*. *Venus*, **7**(4), 173-178 (in Japanese).
- Kanno, S. (1962), Molluscan fauna from the so-called Setana Formation, southwestern Hokkaido, Japan. *Sci. Rept., Tokyo Kyoiku Daigaku, Sec. C.*, **8**(73), 49-62, pls. 1-5.
- Kaufman, Z. S. (1974), On the amphinuclear state of the nucleoli in the oocytes of marine invertebrates. *Zh. Obshch. Biol. (J. general Biol.)*, **35**(1), 144-148 (in Russian).
- Kaufman, Z. S. (1977), *Peculiarities of the sexual cycles of the White Sea invertebrates as an adaptation to the existence in the high latitudes environments. Morphoecological and evolutionary aspects of the problem*. 265 pp. Nauka, Leningrad (in Russian).
- Keen, A. M. (1936), A new pelecypod genus of the family Cardiidae. *Transact. San Diego Soc. Nat. Hist.*, **8**(17), 119-120.
- Keen, A. M. (1937), *An abridged check list and bibliography of West North American marine Mollusca*. 87 pp. Stanford Univ. Press, Stanford, CA.
- Keen, A. M. (1954), Five new species and a new subgenus in the pelecypod family Cardiidae. *Bull. Amer. Paleontol.*, **35**(153), 1-24 (311-330), pl. 1(29).
- Kira, T. (1954 [repr. 1954, 1955, 1956, 1957, 1958]), *Coloured illustrations of the shells of Japan*. 204 pp., 67 pls. Hoikusha, Osaka (in Japanese).
- Kira T. 1959. *Coloured illustrations of the shells of Japan*. Enlarged and revised edition. 239 pp., 71 pls. Hoikusha, Osaka (in Japanese).
- Kira, T. (1962), *Shells of the Western Pacific in color*. 224 pp., 72 pls. Hoikusha, Osaka.
- Klimova, V. L. (1974), Bottom biocenoses of the Peter the Great Bay shelf. In, *Hydrobiology and biogeography of the shelf zones in cold and temperate waters of the World Ocean*. 76. Nauka, Leningrad (in Russian).
- Knipowitsch, N. (1902), Zoologische Ergebnisse der Russischen Expeditionen nach Spitzbergen. Mollusca und Brachiopoda. I. *Annu. Zool. Mus., Imper. Acad. Sci. St.-Petersburg*, **6**, 435-558, Tabs. 18, 19.
- Knipowitsch, N. (1903), Zoologische Ergebnisse der Russischen Expeditionen nach Spitzbergen. Mollusca und Brachiopoda. II, III. *Annu. Zool. Mus., Imper. Acad. Sci. St.-Petersburg*, **7**, 355-459, Tabs. 8-11.
- Kobelt, W. (1888), *Prodromus faunae molluscorum testaceorum maria europaea inhabitantium*. 550 pp. Bauer u. Raspe, Nürnberg.
- Kotaka, T. (1950), A new *Clinocardium* from Aomori Prefecture. *Short Paps., Inst. Geol. Paleontol., Tohoku Univ.*, no. 2, 46-50, pl. 5.
- Kotaka, T. (1962), Marine Mollusca dredged by the "S. S. Hokuho-maru" during 1959 in the Okhotsk Sea. *Sci. Repts. Tohoku Univ.*, 2nd Ser., spec. vol. no. 5, 127-158, pls. 33-35.
- Kotaka, T. and Noda, H. (1967), Miocene Mollusca from the Minami-Tsugaru district, Aomori Prefecture, Northeast Japan. *Res. Bull. Saito Ho-on Kai Mus.*, no. 36, 33-47, pls. 1-2.
- Krause, A. (1885), Ein Beitrag zur Kenntniss der Mollusken-Fauna des Beringsmeeres. Brachiopoda et Lamellibranchiata. *Arch. Naturgesch.*, **1**, 14-40, Taf. 3.
- Kulichkova, V. A. (1955), Feeding of the king crab at shores of Kamchatka and Sakhalin during spring-and-summer season. *Proc. Pac. Res. Inst. Fish. Oceanogr.*, **43**, 21-42 (in Russian).
- Kun, M. S. and Mikulich, L. V. (1954), Structure of food of the Far East commercial crabs during a summer season. *Proc. Pacific Res. Inst. Fish. Oceanogr. (TINRO)*, **41**, 319-322 (in Russian).
- Kuroda, T. (1931). Fossil Mollusca. In Homma, F., *Geology of Central Shinano*. Pt. 4. 1-90, pls. 1-13 (in Japanese).
- Kuroda, T. and Habe, T. (1952), *Check list and bibliography of the Recent marine Mollusca of Japan*. 210 pp. L.W.Stach, Tokyo.
- Kuroda, T. and Kinoshita, T. (1951), A catalogue of marine molluscan shells of Hokkaido. (Icones of marine animals and plants of Hokkaido. Mollusca No. 1). *Bull. Hokkaido Reg. Fish. Res. Lab., Fish. Agency*, no. 2, 1-40 (in Japanese).
- Kuroda, T. and Koba, K. (1933), Molluscan fauna of the Northern Kurile Islands. *Bull. Biogeogr. Soc. Japan*, **4**(2), 151-170, pl. 14 (in Japanese).
- Kuznetsov, A. P. (1961), Materials on the quantitative account of bottom fauna in Kamchatsky Bay. *Transact. Inst. Oceanol., Acad. Sci. USSR*, **46**, 103-123 (in Russian).
- Kuznetsov, A. P. (1963), *Fauna of bottom invertebrates of the Pacific coasts of Kamchatka and northern Kurile Islands*. 271 pp. Acad. Sci. USSR Publ. House, Moscow (in Russian).
- Kuznetsov, A. P. (1970), Conformities of distribution of food groupings of bottom invertebrates in Barents Sea. *Transact. Inst. Oceanol., Acad. Sci. USSR*, **88**, 5-80 (in Russian).
- Kuznetsov, V. V. (1960), *The White Sea and biological features*



- of its flora and fauna. 322 pp. Acad. Sci. USSR Publ. House, Moscow & Leningrad (in Russian).
- Kuznetsov, V. V. (1961), Biological bases of rational use and increases of trade riches of the White sea. *Proc. Acad. Sci. USSR, Ser. Biol.* no. 4, 608-619 (in Russian).
- Kuznetsov, V. V. (1963), Time and temperature conditions of spawning in marine invertebrates. In, *Materials on complex study of the White sea. Vol. 2.* 32-52. Acad. Sci. USSR Publ. House, Moscow & Leningrad (in Russian).
- Kwon, O. K., Park, G. M. and Lee, J. S. (1993), *Coloured shells of Korea.* 445 pp. Academy Publ. Co., Seoul (in Korean).
- La Rocque, A. (1953), Catalogue of the Recent Mollusca of Canada. *Natl. Mus. Canada, Bull.* **129**, ix + 406 pp.
- Leche, W. (1878), Ofversigt ofver de af Svenska expeditionerna till Novaja Semlja och Jenissej 1875 och 1876 insamlade hafs-mollusker. *Handl. Kongl. Vetensk.- Akad.*, **16**(2), 1-86, tafn. 1-2.
- Leche, W. (1883), Ofversigt ofver de af Vega-expeditionen insamlade arktiska hafsmollusker. I. Lamellibranchiata. In, *Vega-Expeditionens vetenskapliga iakttagelser... utgifna af A. E. Nordenskiöld. 3. Bd.* 433-453, tafn. 32-34. F. & G. Beijers, Stockholm.
- Lisitsyn, A. P. (1961), Conformities of coarse clastics carrying by ice. In Strakhov, N. M., Bezrukov, P. L. and Yablokov, V. S. (eds.), *Modern sediments in seas and oceans.* 232-284. Acad. Sci. USSR Publ. House, Moscow (in Russian).
- Lubinsky, I. (1980), Marine bivalve molluscs of the Canadian Central and Eastern Arctic: faunal composition and zoogeography. *Canad. Bull. Fish. Aquat. Sci.* **207**, 111 pp., 11 pls.
- MacGinitie, G. E. (1955), Distribution and ecology of the marine invertebrates of Point Barrow, Alaska. *Smithson. Miscell. Collects.*, **128**(9), 201 pp., 8 pls.
- MacGinitie, N. (1959), Marine Mollusca of Point Barrow, Alaska. *Proc. U. S. Natl. Mus.*, **109**(3412), 59-208, pls. 1-27.
- Madsen, F. J. (1949), Marine Bivalvia. *The Zoology of Iceland, Vol. IV*, pt. 63. 116 pp. Ejnar Munksgaard, Copenhagen & Reykjavik.
- Marincovich, L. jr. (1983), Molluscan paleontology, paleoecology, and North Pacific correlations of the Miocene Tachilni Formation, Alaska Peninsula, Alaska. *Bull. Amer. Paleontol.*, **84**(317), 55-155, pls. 12-23.
- Marincovich, L. jr. and Powell, C. L. 1989. Preliminary Tertiary molluscan biostratigraphy of the Alaska Peninsula, southwestern Alaska. *U. S. Geol. Surv. Open-File Report* 89-674, 2 sheets.
- Matsui, S. (1985), Recurrent molluscan associations of the Omma-Mandanji fauna in the Gojome-Oga area, Northeast Honshu. Pt. 2. Systematic notes on bivalve species. *Transact. Proc. Palaeontol. Soc. Japan, N. S.*, no. 140, 225-239, pls. 31-34.
- McLaughlin, P. A. (1963), Survey of the benthic invertebrate fauna of the Eastern Bering Sea. *U. S. Fish a. Wildlife Service Spec. Sci. Rept.- Fisheries*, no. 401, 1-75.
- McLean, R. A. (1939), The Cardiidae of the Western Atlantic. *Mems. Soc. Cubana Hist. Natur.*, **13**(3), 157-173, pls. 23-26.
- Merklin, R. L., Petrov, O. M., Amitrov, O. V. (1962), *Atlas and guidebook of molluscs of the Chukotsk Peninsula Quaternary deposits.* 83 pp., 12 pls. Publ. by USSR Academy of Sciences, Moscow (in Russian).
- Merklin, R. L., Zarkhidze, V. S. and Ilyina, A. P. (1979), Guidebook of the marine Pliocene-Pleistocene molluscs of the North East of the European part of the USSR. *Transact. Paleontol. Inst., Acad. Sci. USSR*, **173**, 1-96, pls. 1-13 (in Russian).
- Messjatzaw, I. I. (1930), Trawl und Bodengreifer als Fangwerkzeuge der Bodentiere. *Arch. Hydrobiol. Planktonkunde*, **21**(1), 131-146.
- Messjatzaw, I. I. (1931), Molluscs of the Barents Sea. *Transact. State Oceanogr. Inst.*, **1**(1), 3-168 (in Russian).
- Middendorff, A. T. von. (1849), Beitrage zu einer Malacozoologia Rossica. III. Aufzählung und Beschreibung der zur Meeresfauna Russlands gehorigen Zweischaler. *Mems. de l'Acad. Imper. Sci. St.-Petersbourg. Sci. natur.* **6**, 1-94, Taf. 11-21.
- Mighels, J. W. (1843), Catalogue of the marine, fluviatile, and terrestrial shells of the State of Maine, and adjacent ocean. Boston *J. Nat. Hist.*, **4**(3), 308-350.
- Miloslavskaya, N. M. (1958a), Features of benthos distribution and opportunity of its use by cod fishes on East Murman coast. In, *Conformity of congestions and migrations of trade fishes in the Murman coastal zone and their connection with biological, hydrological and hydrochemical processes.* 103-125. Acad. Sci. of the USSR Publ. House, Moscow & Leningrad (in Russian).
- Miloslavskaya, N. M. (1958b), The temperature factor in distribution of bivalve molluscs in East Murman coast. *Transact. Murman Biol. Sta.*, **4**, 140-150 (in Russian).
- Mohr, N. (1786), *Forsøg til en islandsk naturhistorie, med adskillige oekonomiske samt andre anmaerkninger.* xvi + 414 pp., 7 pls. Holm, Copenhagen.
- Mörch, O. A. L. (1870), Catalogue des mollusques du Spitzberg [sic] recueillis par le Dt. H. Kroyer pendant le voyage de la corvette La Recherche en juin 1838. *Méms. Soc. Malacol. Belg.*, **4**, 7-32.
- Morris, P. A. (1952), *A field guide to shells of the Pacific Coast and Hawaii.* xx + 220 pp., 48 pls. Houghton Mifflin, Boston.
- Naumov, A. D. and Fedyakov, V. V. (1990), Bivalve molluscs of the Novosibirsk Islands area. In Golikov, A. N. (ed.), *Ecosystems of Novosibirsk shallow water and fauna of the Laptev Sea and adjacent waters.* 385-410. Nauka, Leningrad (in Russian).
- Naumov, A. D., Scarlato, O. A. and Fedyakov, V. V. (1987), Class Bivalvia Linnaeus, 1758. In Starobogatov, Ya. I. and Naumov, A. D. (eds.), *Molluscs of the White Sea.* 205-257. Nauka, Leningrad (in Russian).
- Neyman, A. A. (1960), Quantitative distribution of benthos in eastern part of Bering Sea. *Zool. Zh. [Zool. J.]*, **39**(9), 1281-1292 (in Russian).
- Noda, H. (1973), Additional notes on the Miyagian marine fauna from the Gobanshoyama Formation, Ojika Peninsula, Miyagi Prefecture, Japan. *Saito Ho-on Kai Mus. Res. Bull.*, no. 42, 31-45, pls. 4, 5.

- Nordsieck, F. (1969), *Die europäischen Meeresmuscheln (Bivalvia). Vom Eismeer bis Kapverden, Mittelmeer und Schwarzes Meer.* 256 SS., 25 Tafn. Gustav Fischer, Stuttgart.
- Ockelmann, W. K. (1958), Marine Lamellibranchiata. *Medd. Grønland*, **122** (4), 1-256, pls. 1-3.
- Odhner, N. (1910), Marine Mollusca of Iceland in the collections of the Swedish State Museum. *Ark. Zool.*, **7**(4), 1-31, pl. 1.
- Odhner, N. H. (1915), Die Molluskenfauna des Eisfjordes. Zoologische Ergebnisse der Schwedischen Expedition nach Spitzbergen 1908 unter Leitung von Prof. G. De Geer, Teil II, 1. *Handl. Kungl. Svensk. Vetenskapsakad.*, **54**(1), 1-274, Taf. 1-13.
- Official index of rejected and invalid works in zoological nomenclature. First installment: names 1-58. London, 1958, 14 pp.
- Ogasawara, K. (1977), Paleontological analysis of Omma fauna from Toyama-Ishikawa area, Hokuriku Province, Japan. *Sci. Repts. Tohoku Univ.*, 2nd Ser., **47**(2), 43-156, pls. 3-22.
- Ogasawara, K., Masuda, K. and Matoba, Y. (editors) (1986), *Neogene and Quaternary molluscs from the Akita oil-field, Japan.* Commemorative Association of Professor Taisuke Takayasu's Retirement and Supporters' Foundation of Mineral Industry Museum, Mining College, Akita University, 310 pp., 85 pls. (in Japanese).
- Oldroyd, I. S. (1925, not "1924" as stated on title page), The marine shells of the west coast of North America, Vol. 1 [Bivalvia]. *Stanford University, Publications, University Series*, Geological Sciences **1**(1), 247 pp., 57 pls.
- Óskarsson, J. (1952), *Skeldýrafána Íslands. I. Samlokur í sjó (Lamellibranchia).* 119 pls. Atvinnudeild háskólans fiskideild, Reykjavík.
- Packard, A. S. (1867), Observations on the glacial phenomena of Labrador and Maine, with a view of the Recent invertebrate fauna of Labrador. *Mems. Boston Soc. Nat. Hist.*, **1**(2), 210-303, pls. 7, 8.
- Petersen, G. Høpner (1968), Marine Lamellibranchiata. In Jensen, A. S., Lundbeck, W. and Mortensen, Th. (eds.), *Zoology of the Faroes.* 1-80. Vald. Petersens, Copenhagen.
- Petersen, G. Høpner (1978), Life cycles and population dynamics of marine benthic bivalves from Disko Bugt area of west Greenland. *Ophelia*, **17**(1), 95-120.
- Petersen, G. Høpner and Kristensen, R. M. (1991), The marine macrofauna at Igloodik, N. W. T. Arctic Canada (not including Polychaeta). In Jørgensen, M. (ed.), *Arctic Biology Course 1989 Igloodik N. W. T. Canada, 1987-91.* Zoological Museum, University of Copenhagen.
- Petrov, O. M. (1966), Stratigraphy and fauna of marine molluscs in the Quaternary deposits of the Chukotsk Peninsula. *Transact. Geol. Inst., Acad. Sci. USSR*, **155**, 1-290, pls. 1-23 (in Russian).
- Petrov, O. M. (1982), Marine molluscs of the Anthropogene from the Northern region of the Pacific. *Transact. Geol. Inst., Acad. Sci. USSR*, **357**, 1-144, pls. 1-24 (in Russian).
- Pfeffer, G. (1886), Übersicht der im Jahre 1881 vom Grafen Waldburg-Zeit im Karischen Meere gesammelten Mollusken. *Abhandl. Naturwissensch. Verein. Hamburg*, **9**(1), 29-40, 1 Taf.
- Pirozhnikov, P. L. (1937), Class Lamellibranchiata - lamellibranch molluscs of the northern seas. In Gaevskaya, N. S. (ed.), *A field-guide to fauna and flora of the northern seas of the USSR*, 74-87, pls. 34-38. Pishchepromizdat, Moscow & Leningrad (in Russian).
- Popova, N. M. (1948), Productivity of the Kara Sea bottom and ways of its use. *Problemy Arktiki [Problems of Arctic]*, no. 2, 76-92 (in Russian).
- Posselt, H. J. (1895), Østgrønlandske mollusker. *Medd. Grønland*, **19**, 61-93, tav. 1.
- Posselt, H. J. and Jensen, Ad. S. (1898), Grønlands brachiopoder og bløddyr. *Conspectus Faunae Groenlandicae. Brachiopoda et Mollusca. Medd. Grønland*, **23**, I-xix, 1-298, tav. 1, 2.
- Powles, P. M. (1965), Life history and ecology of American plaice (*Hyppoglossoides platessoides*) in the Magdalen shallows. *J. Fish. Res. Bd. Canada*, **2**(2), 119-127.
- Reeve, L. A. (1844-1845), *Monograph of the genus Cardium.* In Reeve, L. A. (ed.), *Conchologia iconica: or, illustrations of the shells of molluscous animals.* Vol. 2. 22 pls. (pls. 1-4: Oct. 1844; pls. 5-8: Nov. 1844; pls. 9-12: Dec. 1844; pls. 13-16: Jan. 1845; pls. 17-22: March 1845). Reeve Brothers, London.
- Richards, H. G. (1962), Studies on the marine Pleistocene. Part 1. The marine Pleistocene of the Americas and Europe. Part 2. The marine Pleistocene mollusks of eastern North America. *Transact. Amer. Philos. Soc., New Ser.*, **52**(3), 1-141, pls. 1-21.
- Rogers, J. E. (1920), *The shell book: a popular guide to a knowledge of the families of living mollusks, and an aid to the identification of shells native and foreign.* 485 pp. Doubleday, Page & Co., New York.
- Römer, E. (1868-1869), Die Familie der Herzmuscheln, Cardacea. In *Systematisches Conchilien - Cabinet von Martini und Chemnitz.* Bd. 10, Abtheilung 2. 124 SS., 14 Tfn. (SS. 1-24, Tfn. 1-7, 1868; SS. 25-124, Tfn. 8-14, 1869). Verlag von Bauer und Raspe, Nürnberg.
- Sars, G. O. (1878), *Bidrag til kundskaben om Norges arktiske fauna. I. Mollusca regionis Arcticae Norvegiae. Oversigt over de i Norges arktiske region forekommende bløddyr.* xiii+466, tafn. 1-34, I-XVIII. A.W. Brøgger, Christiania.
- Savilov, A. I. (1961), The ecological characteristic of bottom invertebrate communities of the Sea of Okhotsk. *Transact. Inst. Oceanol., Acad. Sci. USSR*, **46**, 3-84 (in Russian).
- Scarlato, O. A. (1955), Class Bivalve molluscs - Bivalvia. In Ushakov, P. V. (ed.), *Atlas of invertebrates from Far East seas of the USSR.* 185-198, pls. 49-53. Publ. by Acad. Sci. USSR, Moscow & Leningrad (in Russian).
- Scarlato, O. A. (1981), *Bivalve molluscs of the temperate latitudes of the western part of the Pacific Ocean.* 400 pp., 64 pls. Nauka, Leningrad (in Russian).
- Schlesch, H. (1924), Zur Kenntnis der pliocänen Cragformation von Hallbjarnarstadur, Tjörnes, Nordisland und ihrer Molluskenfauna. *Abhandl. Arch. Molluskenk.*, **1**(3), 309-370, Tafn. 1-12.
- Shimamoto, M. and Koike, T. (1986), The molluscan assemblage from the Tentokuji formation, southwest of Mt. Taihei,

- Akita Prefecture. *Saito Ho-on Kai Mus. Nat. Hist. Res. Bull.*, no. 54, 27-47, pls. 4-6.
- Skalkin, V. A. (1963), Nutrition of flounders in southeastern portion of the Bering Sea. *Transact. All-Union Res. Inst. Fish. Oceanogr. (VNIRO)*, **48**, 223-237 (in Russian).
- Slodkewitsch, W. S. (1938), *Tertiary Pelecypoda from the Far East*. The USSR Academy of Sciences Press, Moscow & Leningrad. Pt. 1. Paleontology of USSR, vol. 10, pt. 3, fasc. 18, 508 pp. (in Russian); Pt. 2 Paleontology of USSR, vol. 10, pt. 3, fasc. 19, 275 pp., 104 pls. (in Russian and English).
- Smith, E. A. (1877), On the Mollusca collected during the Arctic Expedition of 1875-76. *Annls. Mag. Nat. Hist.*, 4<sup>th</sup> Ser., **20**, 131-146.
- Soot-Ryen, T. (1932), Pelecypoda with a discussion of possible migrations of Arctic pelecypods in Tertiary times. *Meddr. Zool. Mus. Oslo*, no. 23, 1-35, pls. 1, 2.
- Soot-Ryen, T. (1958), Pelecypods from East-Greenland. *Skrift. Norsk Polarinst.*, no. 113, 1-32.
- Sowerby, G. B., II. (1840 [1841a]), On some new species of the genus *Cardium*, chiefly from the collection of H. Cuming, Esq. *Proc. Zool. Soc. London*, **8**(92), 105-111.
- Sowerby, G. B., II (1841b), *Cardium*. A catalogue of Recent species. In Sowerby, G. B. I and Sowerby, G. B. II., *The conchological illustrations; or coloured figures of all the hitherto unfigured Recent shells, etc.* 8 pp., 71 figs. Sowerby, London.
- Spärck, R. (1933), Contribution to the animal ecology of the Franz Joseph Fiord and adjacent East Greenland waters. *Medd. Grønland*, **1C**, 1-38.
- Spärck, R. (1936), On the relation between metabolism and temperature in some marine lamellibranchs, and its ecological and zoogeographical importance. *Vidensk. Selskab. Biol. Meddr.*, **13**(5), 1-27.
- Spärck, R. (1937), The benthonic animal communities of the coastal waters. *The Zoology of Iceland, Vol. 1, pt. 6.* 45 pp., Levin and Munksgaard, Copenhagen & Reykjavik.
- Stimpson, W. (1851), *Shells of New England. A revision of the synonymy of the testaceous mollusks of New England; with notes on their structure, and their geographical and bathymetrical distribution.* 58 pp., 2 pls. Boston.
- Stimpson, W. (1854), Synopsis of the marine Invertebrata of Grand Manan, or the region about the mouth of the Bay of Fundy, New Brunswick. *Smithson. Contribs. Knowledge*, **6**, 1-68, pls. 1-3.
- Stimpson, W. (1863), Description of a new *Cardium* from the Pleistocene of Hudson's Bay. *Proc. Acad. Nat. Sci. Phila.*, **14**, 58-59.
- Stimpson, W. (1864), Synopsis of the marine Invertebrata collected by the late Arctic Expedition, under Dr. J. J. Hayes. *Proc. Acad. Nat. Sci. Phila.*, **15**, 138-142.
- Thiele, J. (1928), Arktische Loricaten, Gastropoden, Scaphopoden und Bivalven. In *Fauna Arctica. Eine Zusammenstellung der arktischen Tierformen mit besonderer Berücksichtigung des Spitzbergen-Gebietes auf grund der Ergebnisse der Deutschen Expedition in das Nördliche Eismeer im Jahre 1898.* 5. Band, 2. Lief. 563-632, Taf. 10. Gustav Fischer, Jena.
- Thorson, G. (1933), Investigations on shallow water animal communities in the Franz Joseph Fjord (East Greenland) and adjacent waters. *Medd. Grønland*, **100**(2), 1-68, pls. 1-3.
- Thorson, G. (1934), Contributions to the animal ecology of the Scoresby Sound Fjord complex (East Greenland). Hydrography by H. Ussing. *Medd. Grønland*, **100**(3), 1-68, pls. 1-6.
- Thorson, G. (1936), The larval development, growth, and metabolism of Arctic marine bottom invertebrates compared with those of other seas. *Medd. Grønland*, **100**(6), 1-155.
- Thorson, G. (1951), Scaphopoda, Placophora, Solenogastres, Gastropoda Prosobranchiata, Lamellibranchiata. The Godthaab Expedition 1928. *Medd. Grønland*, **81**(2), 1-117.
- Thorson, G. (1957), Bottom communities (sublittoral or shallow shelf). In: Treatise on marine *Ecology*: and paleoecology, vol. 1. *Mem. Geol. Soc. Amer.*, **67**, 461-534.
- Tryon, G. W., jr. (1873), *American marine conchology, etc.* 208 pp. Tryon, G. W., Philadelphia.
- Tseeb, R. Ya. (1964), *A feed and food adaptations in Barents Sea haddock.* 135 pp. Nauka, Moscow & Leningrad (in Russian).
- Tsikhon-Lukanina, E. A. (1987), *Trophology of aquatic molluscs.* 176 pp. Nauka, Moscow (in Russian).
- Turpaeva, E. P. (1948), A feed of some bottom invertebrates of the Barents Sea. *Zhool. Zh. [Zool. J.]*, **27**(6), 503-512 (in Russian).
- Turpaeva, E. P. (1953), A feed and food groupings of marine bottom invertebrates. *Transact. Inst. Oceanol., Acad. Sci. USSR*, **7**, 259-299 (in Russian).
- Uozumi, S. (1962), Neogene molluscan faunas in Hokkaido. Pt. 1. Sequence and distribution of Neogene molluscan faunas. *J. Fac. Sci., Hokkaido Univ.*, Ser. 4, **11**(3), 507-544.
- Ushakov, P. V. (1953 [post - November 5; published under the griff "top secret"; data of the opening for public use is October 20, 1957]), *Fauna of the Sea of Okhotsk and conditions of their existence.* 459 pp. Publ. by Acad. Sci. USSR, [Moscow & Leningrad] (in Russian).
- Vibe, C. (1939), Preliminary investigations on shallow water animal communities in the Upernavik- and Thule-districts (northwest Greenland). *Medd. Grønland*, **124**(2), 1-42.
- Vibe, C. (1950), The marine mammals and the marine fauna in the Thule district (northwest Greenland) with observations on ice conditions in 1939-41. *Medd. Grønland*, **150**(6), 1-115.
- Vinogradov, L. G. and Neyman, A. A. (1969), The bottom population of a shelf of east part of the Sea of Okhotsk and some biological features of king crab. *Okeanologiya [Oceanology]*, **9**(2), 329-340 (in Russian).
- Vinogradova, N. G. (1954), Materials on the quantitative account of bottom fauna in some bays of Okhotsk and Bering Seas. *Transact. Inst. Oceanol., Acad. Sci. USSR*, **9**, 136-158 (in Russian).
- Volova, G. N. and Scarlato O. A. (1980), *Bivalve molluscs of the Peter the Great Bay.* 96 pp. Far East Book Publishing Co., Vladivostok (in Russian).
- Volova, G. N. and Scarlato, O. A. (1991), Fauna and distribution of bivalve molluscs in the north and west portions of the Sea of Okhotsk. In Ivankov, V. N. and Kudryashov, V. A. (eds.), *Biology of fishes and invertebrates of the northern part*


- of Pacific Ocean. 142-181. Far East State Univ. Press, Vladivostok (in Russian).
- Voskuil, R. (1989), De recente Europese Cardiidae. *Vita marina, Zeebiol. Dokument.*, **Juli-okt.**, Tweekleppigen, 247-278.
- Whiteaves, J. F. (1901), Catalogue of the marine Invertebrata of Eastern Canada. *Bull. Geol. Surv. Canada*, no. 722, 1-271.
- Wood, W. (1825), *Index testaceologicus; or a catalogue of shells, British and foreign, arranged according to the Linnean system, etc.* 188 pp., 76 pls. W. Wood, London.
- Yokoyama, M. (1926), Fossil shells from Sado. *J. Fac. Sci., The Imper. Univ. Tokyo*, Sect. 2, **1(8)**, 249-312, pls. 32-37.
- Zatsepin, V. I. (1962), Faunal communities of bottom invertebrates of Murman coast of the Barents Sea and their connection with North Atlantic communities. *Transact. All-Union Hydrobiol. Soc.*, **12**, 245-344 (in Russian).
- Zatsepin, V. I. and Filatova, Z. A. (1961), Bivalve mollusc *Cyprina islandica* (L.), its geographical distribution and significance in bottom animal communities. *Transact. Inst. Oceanol., Acad. Sci. USSR*, **46**, 201-216 (in Russian).
- Zenkevich, L. A. (1947), *Fauna and biological productivity of a sea. Vol. 2. Seas of the USSR.* 588 pp. Sovetskaya Nauka, Moscow (in Russian).
- Zenkevich, L. A. (1951), *Fauna and biological productivity of a sea. Vol. 1. World Ocean.* 506 pp. Sovetskaya Nauka, Moscow (in Russian).
- Zenkevich, L. A. (1963). *Biology of seas of the USSR.* 739 pp. Acad. Sci. USSR. Publ. House, Moscow (in Russian).
- Zenkevich, L. A. and Brotskaya, V. A. (1937), Materials on ecology of the managing forms of the Barents Sea benthos. *Mems. Moscow State Univ.*, Issue 13, Zoology, 203-226.
- Zhidkova, L. S., Kusina, I. N., Lautenschlager, F. G. and Popova, L. A. (1968), *Atlas of the Upper Miocene and Pliocene molluscs of Sakhalin.* 180 pp., 50 pls. Nauka, Moscow (in Russian).
- Zhidkova, L. S., Bezv, V. E., Ilyina, A. P., Krishtofovich, L. V., Neverova, T. I., Savizky, V. O. and Scheremetjeva, G. N. (1972), *Atlas of Neogene molluscs of the Kurile Islands.* 164 pp., 48 pls. Nauka, Moscow (in Russian).
- Zhirmunsky, A. V. (1969), Comparative research of thermal resistance in the White Sea molluscan cells in connection with vertical distribution and history of fauna formation. *J. Gen. Biol. [Zh. Obshch. Biol.]*, **30(6)**, 686-704 (in Russian).
- Zinbo, N. (1973), Fossil Mollusca from the Utsutoge formation, Iide-machi, Nishiokitama-gun, Yamagata Prefecture, Japan. *Sci. Repts. Tohoku Univ.*, Ser. 2, spec. vol. no. 6 (Hatai Mem. Vol.), 157-162, pl. 14.

Manuscript accepted on 25<sup>th</sup> February, 2001

## Explanation of Plate 1

- Figs. 1-11, 18. *Ciliatocardium ciliatum ciliatum* (Fabricius, 1780).  
Lectotype of *Cardium ciliatum* Fabricius, 1780 - ZMK, without reg. no.; left valve exterior (fig. 1) and interior (fig. 2), right valve exterior (fig. 3) and interior (fig. 4). Greenland; Recent.  
Paralectotype of *Cardium ciliatum* Fabricius, 1780 - ZMK, without reg. no.; left valve exterior (fig. 5) and interior (fig. 6), right valve exterior (fig. 7) and interior (fig. 8). Greenland; Recent.  
Possible syntype of *Cardium arcticum* Sowerby, 1841 - BMNH no. 1975618; left valve exterior (fig. 9), right valve exterior (fig. 10) and interior (fig. 11); "Arctic seas"; Recent.  
Original figure reproduction (Chemnitz, 1782, Taf. 19, Fig. 195) of *Cardium islandicum* Bruguière, 1789; right valve exterior (fig. 18); Iceland (?); Recent.
- Fig. 12. *Ciliatocardium ciliatum dawsoni* (Stimpson, 1863).  
Original figure reproduction (Stimpson, 1863: 58, text-fig.) of *Cardium dawsoni* Stimpson, 1863; Cape Hope, on the southeast side of Hudson's Bay, Arctic Canada; Pleistocene (?).
- Figs. 13-17. *Ciliatocardium ciliatum* (Fabricius, 1780).  
Lectotype of *Cardium californiense* var. *comoxense* Dall, 1900 - USNM no. 427772; left (fig. 13) and right (fig. 14) valve exterior, left (fig. 15) and right (fig. 16) valve interior; Vancouver Island, British Columbia, Canada; Pleistocene.  
Holotype of *Ciliatocardium olchovensis* Petrov, 1982 - GIN no. 23/3829; external cast of left valve (fig. 17); Pamyatnyi Stream, Kamchatskiy Peninsula, east Kamchatka, Russia; lower sub-suite of Ol'khovskaya Suite, Eopleistocene.

All figures in natural size.



## Plate 2

Figs. 1-13.

*Ciliatocardium ciliatum ciliatum* (Fabricius, 1780).


Iceland; ZIsp - cat. no. 1; left valve exterior (fig. 1) and interior (fig. 2), right valve exterior (fig. 3) and interior (fig. 4).

White Sea, off Belaya Inlet, depth 14 m, muddy sand; ZIsp - cat. no. 126; left valve exterior (fig. 5) and interior (fig. 6), right valve exterior (fig. 7) and interior (fig. 8).

Barents Sea, 70°58' N 41°25' E, depth 79 m, pebbles; ZIsp - cat. no. 67; right valve exterior (fig. 9) and interior (fig. 10), left valve interior (fig. 11).

Barents Sea, 69°36.5' N 32°13.5' E, depth 162 m; ZIsp - cat. no. 60; right (fig. 12) and left (fig. 13) valves exterior.

All figures in natural size.



### Plate 3

Figs. 1-4. *Ciliatocardium ciliatum nordenskiöldi* Kafanov, 1981.

Holotype - ZIsp no. 1/200156; left valve exterior (fig. 1) and interior (fig. 2), right valve exterior (fig. 3) and interior (fig. 4); Laptev Sea, Komsomol'skoy Pravdy Islands, between Angreya Island and Samuila Island, depth 37 m, mud.

Figs. 5-8. *Ciliatocardium ciliatum ochotense* Kafanov, 1981.


Holotype - ZIsp no. 1/200366; left valve interior (fig. 5) and exterior (fig. 6), right valve exterior (fig. 7) and interior (fig. 8); north coast of Okhotsk Sea, west from Tauyskaya Inlet, depth 121 m, mud.

Figs. 9-12. *Ciliatocardium ciliatum pubescens* (Couthouy, 1838).

Off Newfoundland, east Canada, 43°07.5' N 50°22.5' W, depth 43 m; ZIsp - cat. no. 1; right valve exterior (fig. 9) and interior (fig. 10), left valve exterior (fig. 11) and interior (fig. 12).

All figures in natural size.






## Plate 4

- Figs. 1-4. *Ciliatocardium ciliatum tchuktchense* Kafanov in Scarlato, 1981.  
Chukotka Peninsula, Russia, Provideniya Bight, depth 22-11 m, mud; paratype - ZIsp cat. no. 2; left valve exterior (fig. 1) and interior (fig. 2), right valve interior (fig. 3) and exterior (fig. 4).
- Fig. 5. *Ciliatocardium ciliatum pubescens* (Couthouy, 1838).  
Original figure reproduction (Couthouy, 1838: 61, pl. 3, fig. 6) of *Cardium pubescens* Couthouy, 1838; Massachusetts Bay, Massachusetts, U.S.A.
- Figs. 6-7. *Ciliatocardium ciliatum* cfr. *dawsoni* (Stimpson, 1863).  
Arctic coast of Alaska, 71°31' 20" N 156°23' W, Pleistocene (?); ZIsp no. 1/13466 (former CAS no. 34343); left valve exterior (fig. 6) and interior (fig. 7).
- Figs. 8-9. *Ciliatocardium ciliatum* (Fabricius, 1780).  
Holotype of *Papyridea sertunayanum* Slodkewitsch, 1938 - CNIGRM no. 185/5294; deformed external cast of right valve (fig. 8); between the mouths of Noyami and Malyi Sertunay Rivers, Aleksandrovsk-Sakhalinsky District, Sakhalin, Russia; middle Miocene Aleksandrovskaya and/or Sertunayskaya Suite.  
CNIGRM no. 142/5294; right valve exterior (fig. 9); the same locality as fig. 8.
- Fig. 10. *Ciliatocardium ciliatum ciliatum* (Fabricius, 1780).  
Possible syntype of *Cardium arcticum* Sowerby, 1841 - BMNH no. 1975618; left valve interior; "Arctic seas"; Recent.

All figures in natural size.



## Plate 5

Figs. 1-13.

*Ciliatocardium ciliatum nordenskiöldi* Kafanov, 1981.

Laptev Sea, Komsomol'skoy Pravdy Islands, between Angreya Island and Samuila Island, depth 37 m, mud; ; paratype - ZIsp cat. no. 10; left valve exterior (fig. 1) and interior (fig. 2), right valve exterior (fig. 3) and interior (fig. 4).

Laptev Sea, off Severnaya Zemlya Archipelago, depth 40 m; paratype - ZIsp cat. no. 45; left (fig. 5) and right (fig. 6) valves exterior.

Central part of Barents Sea, depth 65 m; paratype - ZIsp cat. no. 4; left (fig. 7) and right (fig. 8) valves exterior, left (fig. 9) and right (fig. 10) valves interior.

Kara Sea, off entrance of Karskie Vorota Strait; paratype - ZIsp cat. no. 41; right valve exterior (fig. 11) and interior (fig. 12), left valve exterior (fig. 13).

Figs. 14-18.


*Ciliatocardium ciliatum ochotense* Kafanov, 1981.

Okhotsk Sea, Nagaeva Bight; paratype - ZIsp cat. no. 8; right valve exterior (fig. 14).

Okhotsk Sea, Nagaeva Bight; paratype - ZIsp cat. no. 6; right valve exterior (fig. 15) and interior (fig. 16), left valve exterior (fig. 17).

Northwestern coast of Okhotsk Sea; paratype - ZIsp cat. no. 7; left valve exterior (fig. 18).

All figures in natural size.



## Plate 6

Figs. 1-6, 13-15. *Ciliatocardium ciliatum* tchuktchense Kafanov in Scarlato, 1981.

Holotype - Zisp no. 1/200291; left valve exterior (fig. 1) and interior (fig. 2), right valve exterior (fig. 3) and interior (fig. 4).

Bering Sea, Provideniya Bight, depth 22-11 m; paratype - ZIsp cat. no. 43; left (fig. 5) and right (fig. 6) valves exterior.


Bering Sea, Bristol Bay; paratype - ZIsp cat. no. 149; left (fig. 13) and right (fig. 14) valves exterior, right valve interior (fig. 15).

Fig. 7-12. *Ciliatocardium ciliatum pubescens* (Couthouy, 1838).

Off Newfoundland, 44°56.4' N 49°01.7' W, east Canada; ZIsp cat. no. 4; left (fig. 7) and right (fig. 8) valves exterior.

Off Newfoundland, 43°07.5' N 50°22.5' W, east Canada; ZIsp cat. no. 1; left valve exterior (fig. 9) and interior (fig. 10), right valve exterior (fig. 11) and interior (fig. 12).

All figures in natural size.



## Plate 7

- Figs. 1-4. *Ciliatocardium ciliatum tchuktchense* Kafanov in Scarlato, 1981.  
Okhotsk Sea, southern Sakhalin, southwestern part of Terpeniya Bay, depth 40 m; paratype - ZIsp cat. no. 96; left (fig. 1) and right (fig. 2) valves exterior.  
Bering Sea, Anadyrskiy Bay, depth 25 m; paratype - ZIsp cat. no. 43; left (fig. 3) and right (fig. 4) valves exterior.
- Figs. 5-6. *Ciliatocardium ciliatum chikagawaense* (Kotaka, 1950).  
Holotype of *Clinocardium chikagawaense* Kotaka, 1950 - IGPS no. 72999; right valve exterior (fig. 5) and interior (fig. 6); the sea cliff at the outlet of Chikagawa River at Chikagawa, Tanabu-machi, Shimokita District, Aomori Prefecture, Honshu, Japan; Pliocene Hamada Formation.
- Figs. 7-8. *Ciliatocardium ciliatum maruyamense* Kafanov et Savizky, 1982.  
Holotype - PIN no. 28/3962; external cast of left (fig. 7) and right (fig. 8) valves; Malyi Takoy River, Dolinsk District, Sakhalin, Russia; lower sub-suite of Maruyamskaya Suite (stratotypical section), upper Miocene.
- Figs. 9-16. *Ciliatocardium likharevi* Kafanov in Scarlato, 1981.  
Holotype - ZIsp no. 1/200274; left (fig. 9) and right (fig. 10) valves exterior, right (fig. 11) and left (fig. 12) valves interior; Japan Sea, Tatar Strait, off Rakuma (now Antonovo Village), Sakhalin, Russia, depth 45-40 m, sand.  
Off Nagasaki, Kyushu, Japan; paratype - ZIsp cat no. 22/200290; right valve exterior (fig. 13) and interior (fig. 14).  
South Kurile shallow-water, between Kunashir and Shikotan Islands; paratype - ZIsp cat. no. 8; left valve exterior (fig. 15) and interior (fig. 16).

All figures in natural size.



