

A first record of *Pugettia* Dana (Crustacea: Decapoda: Brachyura) from the Pliocene of Japan

Hiroaki Karasawa* and Koushi Kitamura**

*Mizunami Fossil Museum, Yamanouchi, Akeyo, Mizunami,
Gifu 509-6132, Japan <GHA06103@nifty.com>

**Ichino-cho 2396-6, Hamamatsu, Shizuoka 435-0051, Japan

Abstract

A new majoid crab, *Pugettia yamafui*, is described from the upper Pliocene Kakegawa Group in Shizuoka Prefecture, Honshu, Japan. Recognition of this species extends the geologic range for the genus back to the late Pliocene.

Key words: Crustacea, Decapoda, Brachyura, Majoidea, *Pugettia*, Pliocene, Japan

Introduction

The genus *Pugettia* Dana, 1851 (Majoidea: Epialtidae) occurs mostly in the north Pacific, and comprises 12 species from the northwest Pacific, six species from the northeast Pacific, and only one species from the Tasman Sea (Garth, 1958; Griffin & Tranter, 1986; Richer de Forges, 1993). Among these, *Pugettia producta* (Randall, 1839) and *P. richi* Dana, 1851, from the Pleistocene of California (Rathbun, 1926) are known in the fossil record.

The purpose of the present paper is to describe a new species of *Pugettia* from the late Pliocene of Japan. The specimen was collected from sandstone of the Tenno silty sandstone Member of the Dainichi Formation exposed at Kamisaigo, Kakegawa City, Shizuoka Prefecture. *Pugettia* occurred in the deposits, in association with decapods, *Pagurus* spp., *Dardanus* spp., and *Carcinoplax longimanus* (De Haan, 1833) (Kitamura unpubl.). Shiba et al. (2001) indicated that the geologic age of the Dainichi Formation was the latest Pliocene (about 2 Ma) based upon the fission-track dating. Shiba et al. (2001) gave detailed discussions of the Dainichi Formation of the present locality.

The specimen is housed in the Mizunami Fossil Museum, Yamanouchi, Akeyo, Mizunami, Gifu, Japan.

Systematics

Family Epialtidae MacLeay, 1838

Genus *Pugettia* Dana, 1851

Type species: *Pugettia gracilis* Dana, 1851 by subsequent designation of Miers (1879).

Geologic range: L. Pliocene-Recent.

Pugettia yamafui, new species

(Fig. 1)




Fig. 1. *Pugettia yamafui*, new species. MFM39019 (holotype), dorsal view of carapace, $\times 2.0$.

Material examined: MFM39019 (holotype).

Diagnosis: Moderate sized *Pugettia*. Postorbital spine and hepatic lobe fused, forming wing-like plate. Dorsal surface well areolated. Gastric region with four tubercles. Cardiac region with large,

median tubercle. Each epibranchial region with three tubercles set in inverted triangle.

Etymology: Named by an arbitrary combination of letters.

Description: Carapace moderate in size, pyriform, much longer than wide. Rostrum bifid, rostral spines broken. Praeorbital spine prominent, directed forward. Postorbital spine and hepatic lobe fused, forming wing-like plate. Lateral branchial tubercle small, slightly directed posteriorly. Posterior half of posterolateral and posterior margins not preserved. Dorsal surface strongly convex, well areolated. Epigastric ridge weakly developed. Gastric region strongly inflated with four tubercles; two on mid-line, posterior one largest; tubercle on each posterior protogastric region. Cardiac region strongly ridged with large, median tubercle. Intestinal region not preserved. Epibranchial region strongly inflated with three tubercles set in inverted triangle. Each mesobranchial region with short, oblique ridge.

Discussion: *Pugettia yamafui* is most similar to the extant *Pugettia incisa* (De Haan, 1839) among known Recent species by having a fused postorbital spine and hepatic lobe, but differs in that there are four tubercles on the gastric region and three on the epibranchial region. Hitherto, known fossil members of *Pugettia* have been recorded from the Pleistocene of California (Rathbun, 1926). The discovery of the new species extends the geologic range for the genus back to the late Pliocene.

References

- Dana, J. D. (1851), On the classification of the maioid Crustacea or Oxyrhyncha. *Amer. J. Sci. Arts.*, (2), **11**, 425-434.
- Garth, J. S. (1958), Brachyura of the Pacific Coast of America: Oxyrhyncha. *Allan Hancock Pacific Exped.*, **21**, 1-854.
- Griffin, J. G. and H. A. Tranter (1986), The Decapoda Brachyura of the Siboga Expedition. Part VIII Majidae. *Siboga-Expedite Monogr.*, **39c4**, 1-335.
- Haan, W. de (1833-1850), Crustacea. In P. F. von Siebold, *Fauna Japonica sive Descriptio Animalium, quae in Itinere per Japoniam, Jussu et Auspiciis Superiorum, qui Summun in India Batava Imperium Tenent, Suscepto, Annis 1823-1830 Collegit, Notis, Observationibus et Adumbrationibus Illustravit*, Ludguni-Batavorum [Leiden], i-xvii, i-xxxii, ix-xvi, 1-243 pp., pls. A-J, L-Q, 1-55.
- Miers, E. J. (1879), On the classification of the majoid Crustacea or Oxyrhyncha, with a synopsis of the families, subfamilies and genera. *J. Linnean Soc. Lond. (Zool.)*, **14**, 634-673.
- Randall, J. W. (1839), Catalogue of the Crustacea Brought by Thomas Nuttall and J. K. Townsend, from the West Coast of North America and the Sandwich Islands, with Descriptions of Such Species As Are Apparently New, among Which Are Included Species of Different Localities, Previously Existing in the Collection of the Academy. *Jour. Acad. Nat. Sci. Philadelphia*, **8**, 106-147.
- Richer de Forges, B. (1993), Deep sea crabs of the Tasman Seamounts (Crustacea: Decapoda: Brachyura). *Rec. Australian Mus.*, **45**, 11-24.
- Shiba, M., K. Yokoyama, T. Shinmura, and K. Kitamura (2001), Excavation report of Pliocene fossil Cetacea from the Kakegawa Group, Kamisaigo, Kakegawa City, central Japan - Geology and sedimentary environment. *Sci. Rep. Mus. Tokai Univ.*, **3**, 77-89. (in Japanese with English abstract)
- MacLeay, W. S. (1838), On the brachyurous decapod Crustacea brought from the Cape by Dr. Smith. In *Illustrations of the Annulosa of South Africa; being a Portion of the Objects of Natural History Chiefly Collected during an Expedition into the Interior of South Africa, under the Direction of Dr. Andrew Smith, in the Years 1834, 1835, and 1836; Fitted out by "The Cape of Good Hope Association for Exploring Central Africa"*, London. p. 53-71.
- Rathbun, M. J. (1926), The fossil stalk-eyed Crustacea of the Pacific slope of North America. *U.S. Natn. Mus. Bull.*, **138**, 155 p.

Manuscript accepted on 3rd July, 2003